

10/050,704



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : C12N 15/57, 9/64, C07K 16/40, 19/00, C12N 1/21, 5/10	<b>A1</b>	(11) International Publication Number: <b>WO 99/35276</b>  (43) International Publication Date: 15 July 1999 (15.07.99)
(21) International Application Number: PCT/US98/26214  (22) International Filing Date: 16 December 1998 (16.12.98)  (30) Priority Data: 09/005,263      9 January 1998 (09.01.98)      US  (71) Applicant: SCHERING CORPORATION [US/US]; 2000 Gal- loping Hill Road, Kenilworth, NJ 07033-0530 (US).  (72) Inventors: DE SAINT-VIS, Blandine, Marie; 76, rue d'Anvers, F-69007 Lyon (FR). FOSSIEZ, Francois; 111, rue du Berger, F-69280 Marcy l'Etoile (FR). CAUX, Christophe; 1, rue de Savoie, F-69002 Lyon (FR). LEBECQUE, Serge, J., E.; 514, chemin du Marand, F-69380 Civrieux d'Azergue (FR).  (74) Agents: MCLAUGHLIN, Jaye, P. et al.; Schering-Plough Corporation, Patent Dept. K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV, MD, MG, MK, MN, MX, NO, NZ, PL, PT, RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: MAMMALIAN MATRIX-TYPE METALLOPROTEASE

## (57) Abstract

Nucleic acids encoding a membrane type matrix metalloprotease, from a mammal, reagents related thereto, including specific antibodies, and purified proteins are described. Methods of using said reagents and related diagnostic kits are also provided.

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## MAMMALIAN MATRIX-TYPE METALLOPROTEASE

5

This filing claims priority from United States Application Serial No. 09/005,263, filed January 9, 1998, the disclosure of which is hereby fully incorporated by reference.

10

## FIELD OF THE INVENTION

The present invention contemplates compositions related to proteins from animals, e.g., mammals, which function as proteinases. In particular, it provides nucleic acids which encode the proteinases, antibodies to, and proteins which exhibit biological functions, e.g., capacity to degrade proteinaceous substrates.

15

## BACKGROUND OF THE INVENTION

The proteases are a very broad group of enzymes which carry out an enzymatic function of hydrolyzing a peptide bond. See, e.g., Beynon (ed. 1989) Proteolytic Enzymes: A Practical Approach IRL Press, Oxford; Methods in Enzymology vols. 244 and 248. Within the group, there is a wide range of substrate specificities for the amino acids adjacent the cleavage sites. Proteases are typically categorized on the basis of their catalytic mechanisms, e.g., based upon studies of their active sites, or by the effects of pH. Four main categories of proteases are serine proteinases, sulfhydryl proteases, acid proteases, and metalloproteases. They may also be classified according to their sites of substrate cleavage, e.g., endoproteases, amino peptidases, or carboxy peptidases.

20

25

30

35

40

Proteases have traditionally held a large share of the industrial enzyme market. Proteases are used in many industrial processes, including in detergents and cleaning products, e.g., to degrade protein materials such as blood and stains, in leather production, e.g., to remove hair, in baking, e.g., to break down glutens, in

flavorings, e.g., soy sauce, in meat tenderizing, e.g., to break down collagen, in gelatin or food supplement production, in the textile industry, in waste treatment, and in the photographic industry. See, e.g., Gusek  
5 (1991) Inform 1:14-18; Zamost, et al. (1996) J. Industrial Microbiol. 8:71-82; James and Simpson (1996) CRC Critical Reviews in Food Science and Nutrition 36:437-463; Teichgraeber, et al. (1993) Trends in Food Science and Technology 4:145-149; Tjwan, et al. (1993) J.  
10 Dairy Research 60:269-286; Haard (1992) J. Aquatic Food Product Technology 1:17-35; van Dijk (1995) Laundry and Cleaning News 21:32-33; Nolte, et al. (1996) J. Textile Institute 87:212-226; Chikkodi, et al. (1995) Textile Res. J. 65:564-569; and Shih (1993) Poultry Science  
15 72:1617-1620.

Matrix metalloproteinases (MMPs) are a family of enzymes whose main physiological function is degradation of the extracellular matrix. See, e.g., Parsons, et al. (1997) Br. J. Surgery 84:160-166. These enzymes are  
20 present in normal healthy individuals and have been shown to have an important role in processes such as wound healing (see Wolf, et al. (1992) J. Invest. Dermatol. 99:870-872; and Wysocki, et al. (1993) J. Invest. Dermatol. 101:64-68), pregnancy and parturition (see  
25 Jeffrey (1991) Seminars Perinatol. 15:118-126), bone resorption (see Delaisse and Vaes, pp. 290-314 in Rifkin and Gay (eds. 1992) Biology and Physiology of the Osteoclast CRC Press, Ratan, FL), and mammary involution (Talhouk, et al. (1992) J. Cell Biol. 118:1271-1282).  
30 See also Nagase (1996) in Hooper (ed.) Zinc Metalloproteinases in Health and Disease Taylor and Francis, London. A recent focus on the MMPs is on their role in certain disease states in which breakdown of the extracellular matrix is a key feature, e.g., in  
35 rheumatoid arthritis (see Harris (1990) NEJ Med. 322:1277-1289), periodontal disease (see Page (1991) J. Periodont. Res. 26:230-242), and cancer (see Brown (1997) Medical Oncology 14:1-10; Chambers and Matisian (1997) J. NCI 89:1260-1270; Yu, et al. (1997) Drugs and Aging



11:229-244; Yu, et al. (1997) Clinical Pharmacology  
11:229-244; Wojtowicz-Praga, et al. (1997) Invest. New  
Drugs 15:61-75; Coussens and Werb (1996) Chem. Biol.  
3:895-904; and Talbot and Brown (1996) Eur. J. Cancer  
5 32A:2528-2533).

While there are many uses for proteases, there is  
always the need for a more active or specific protease  
under various specific conditions. Alternatively, the  
distribution of these gene products may be useful as  
10 markers for specific cell or tissue types. There is a  
need for new proteinases of differing properties,  
specificities, and activities.

#### SUMMARY OF THE INVENTION

15 In a search for DC restricted molecules, a novel  
member of the MMP family of proteolytic enzymes was  
identified which belongs to the Membrane-type Matrix  
Metalloproteinase (MT-MMP) subclass. This fifth MT-MMP  
proteinase, located on chromosome 16p13.3, is present in  
20 spleen, lymph node, thymus, appendix, PBL, and bone  
marrow, and strongly expressed by DC and weakly by  
granulocytes and effector T cells. Interestingly, the  
mRNA expression of this gene is down-regulated by CD40L  
activation of CD34<sup>+</sup>- and monocyte-derived DC. According  
25 to its cellular expression and putative membrane  
localization, a role is proposed for this novel  
Membrane-type Matrix Metalloproteinase gene in  
degradation of the extracellular matrix during DC  
migration.

30 The present invention provides a binding compound  
comprising an antibody binding site which specifically  
binds to primate F06B09 protein; a nucleic acid  
comprising sequence encoding at least 12 amino acids of  
primate F06B09 protein; a substantially pure protein  
35 which is specifically recognized by the above antibody  
binding site; a substantially pure primate F06B09 protein  
or peptide thereof; and a fusion protein comprising a 30  
amino acid sequence portion of primate F06B09 protein  
sequence.

In certain binding compound embodiments, the antibody binding site is specifically immunoreactive with a protein selected from polypeptides of SEQ ID NO: 4; is raised against a purified or recombinantly produced  
5 primate F06B09 protein; is immunoselected on a substantially purified or recombinantly produced primate F06B09 protein; is in a monoclonal antibody, Fab, or F(ab)2; is detectably labeled; is attached to a solid  
10 substrate; is from a rabbit or mouse; binds with a Kd of at least about 300  $\mu$ M; is fused to another protein segment; is in a chimeric antibody; or is coupled to another chemical moiety.

The invention also provides a method of making an antigen-antibody complex, comprising a step of contacting  
15 a primate biological sample to a specific binding antibody described. In preferred embodiments, the method further includes steps to purify the antigen or antibody.

Alternative embodiments provide an antibody binding site wherein the binding site is detected in a biological  
20 sample by a method comprising the steps of contacting a binding agent having an affinity for F06B09 protein with the biological sample; incubating the binding agent with the biological sample to form a binding agent:F06B09 protein complex; and detecting the complex. In certain  
25 embodiments, the biological sample is human, and the binding agent is an antibody.

The invention also provides kits containing a composition described above and instructional material for the use of the composition; or segregation of the  
30 composition into a container. Typically, the kit is used to make a qualitative or quantitative analysis.

The invention also embraces a cell comprising an antibody described above; a cell transfected with a nucleic acid described above; or a cell comprising a  
35 fusion protein described above.

In nucleic acid embodiments, the nucleic acid may encode a polypeptide which specifically binds an antibody generated against an immunogen selected from the group consisting of the mature polypeptides of SEQ ID NO: 4.

Alternatively, the nucleic acid may encode at least 12 amino acids of SEQ ID NO: 4; comprise sequence of at least about 39 nucleotides selected from protein coding portions of SEQ ID NO: 1 or 3; hybridize to SEQ ID NO: 1 or 3 under stringent wash conditions of at least 45° C and less than about 150 mM salt; comprise sequence made by a synthetic method; be an expression vector; be detectably labeled; be attached to a solid substrate; be from human; bind with a Kd of at least about 300  $\mu$ M; be fused to another nucleic acid segment; be coupled to another chemical moiety; be operably associated with promoter, ribosome binding site, or poly-A addition site; be a PCR product; be transformed into a cell, including a bacterial cell; be in a sterile composition; be capable of selectively hybridizing to a nucleic acid encoding an F06B09 protein; comprise a natural sequence; comprise a mature protein coding segment of SEQ ID NO: 1 or 3; encode proteolytically active portion of F06B09; be detected in a biological sample by a method comprising: contacting a biological sample with a nucleic acid probe capable of selectively hybridizing to said nucleic acid, incubating the nucleic acid probe with the biological sample to form a hybrid of the nucleic acid probe with complementary nucleic acid sequences present in the biological sample; and determining the extent of hybridization of the nucleic acid probe to the complementary nucleic acid sequences, including the method where the nucleic acid probe is capable of hybridizing to a nucleic acid encoding a protein selected from the group consisting of the mature polypeptides of SEQ ID NO 4.

In protein or polypeptide embodiments, the proteins may bind with a Kd of at least about 300  $\mu$ M to an antibody generated against an immunogen of the polypeptides of SEQ ID NO: 4; be immunoselected on an antibody which selectively binds a polypeptide of SEQ ID NO: 4; comprise sequence of at least 12 contiguous residues of SEQ ID NO: 4; exhibit a post-translational modification pattern distinct from natural F06B09; be 3-

fold or fewer substituted from natural sequence; be recombinantly produced; be denatured; have sequence of full length natural polypeptide; be detectably labeled; be attached to a solid substrate; be from human; be in a sterile composition; be fused to another protein segment; be coupled to another chemical moiety; comprise at least a fragment of at least 32 amino acid residues from a human F06B09 protein; comprise mature polypeptide sequence selected from the group consisting of SEQ ID NO 2 and 4; be a soluble protein; be a naturally occurring protein; or be a proteolytically active portion of F06B09.

The invention also provides an isolated protein which specifically binds to an antibody generated against an immunogen selected from the group consisting of the full length polypeptides of SEQ ID NO: 4. Preferably such protein binds to the antibody with a  $K_d$  of at least about 300  $\mu M$ ; is immunoselected on an antibody which selectively binds a polypeptide of SEQ ID NO: 4; comprises sequence of at least 12 contiguous residues of SEQ ID NO: 4; exhibits a post-translational modification pattern distinct from natural F06B09; is 3-fold or fewer substituted from natural sequence; is recombinantly produced; is denatured; has sequence of full length natural polypeptide; is detectably labeled; is attached to a solid substrate; is from human; is in a sterile composition; is fused to another protein segment; is coupled to another chemical moiety; comprises at least a fragment of at least 32 amino acid residues from a human F06B09 protein; comprises mature polypeptide sequence selected from the group consisting of SEQ ID NO 4; is a soluble protein; or comprises a proteolytic activity of F06B09.

In certain other embodiments, the invention embraces a fusion protein described above, which comprises sequence from an enzymatically active portion of SEQ ID NO: 4. Preferably such protein binds with a  $K_d$  of at least about 300  $\mu M$  to an antibody generated against an immunogen having sequence of a polypeptide of SEQ ID NO.

4; is immunoselected on an antibody which selectively binds a polypeptide of SEQ ID NO: 4; comprises sequence of at least 12 contiguous residues of SEQ ID NO: 4; is recombinantly produced; is denatured; has sequence of full length natural polypeptide; is detectably labeled; is attached to a solid substrate; comprises sequence from human; is in a sterile composition; is fused to another protein segment; is coupled to another chemical moiety; comprises at least a fragment of at least 32 amino acid residues from a human F06B09 protein; comprises mature polypeptide sequence from SEQ ID NO 4; is a soluble protein; or comprises a proteolytic activity of F06B09.

The invention also provides a substantially pure protein described above which comprises a proteolytic activity of F06B09.

A method of modulating physiology or development of a cell comprising contacting said cell with said compositions is provided.

20

## DETAILED DESCRIPTION

## I. General.

Dendritic cells (DC), present in all lymphoid and non lymphoid organs, are professional antigen presenting cells (APC) which have the unique capacity to activate naive T cells. See, e.g., Banchereau and Steinman (1998) Nature 392:245-252; and Steinman (1991) Annu. Rev. Immunol. 9:271-296. DC, originated from bone-marrow, migrate as precursors through bloodstream to non lymphoid tissues where, at immature stage, DC such as the epidermal Langerhans cells capture antigens with high efficiency and become circulating veiled cells. These cells bearing antigens migrate from the peripheral non lymphoid tissues via lymphatics or bloodstream into lymphoid tissues where they localize in T cell-rich areas as mature interdigitating DC (IDC). See, e.g., Austyn (1996) J. Exp. Med. 183:1287-1292; Austyn, et al. (1988) J. Exp. Med. 167:646-651; Fossum (1988) Scand. J. Immunol. 27:97-105; Hoefsmit, et al. (1982) Immunobiology. 161:255-265; Kripke, et al. (1990) J.

Immunol. 145:2833-2838; Larsen, et al. (1990) J. Exp. Med. 172:1483-1494; Macatonia, et al. (1987) J. Exp. Med. 166:1654-1667; Romani, et al. (1989) J. Exp. Med. 169:1169-1178. At this site, IDC efficiently present  
5 processed Ags to naive T cells and generate a specific immune response. See, e.g., Inaba, et al. (1983) Proc. Natl. Acad. Sci. USA. 80:6041-6045; and Inaba and Steinman (1985) Science. 229:475-479. Thus, migration constitutes an integral part of DC function.

10 The recruitment of DC into a site of tissue damage and the subsequent migration of DC into secondary lymphoid organs is dependent upon a dynamic and complex series of events, including activation by inflammatory stimuli. See, e.g., Butcher (1991) Cell. 67:1033-1036.

15 This mechanism, implies transendothelial migration beyond the vascular compartment involving the expression of integrin molecules, the movement along leukocyte specific chemotactic gradients (Taub (1996) Cytokine Growth Factor Rev. 7:355-376) and possibly the secretion of matrix-degrading enzymes (Watanabe, et al. (1993) J. Cell Sci. 20 104:991-999). In addition, the trafficking of DC into tissues involves breaching the basement membrane (dermo-epidermic junction), which would necessitate the production of a matrix-degrading enzyme.

25 Matrix metalloproteinases, or matrixins, represent a group of structurally related zinc-dependent endopeptidases that are involved in extracellular matrix and basement membrane degradation and cell-matrix interactions. See, e.g., Basbaum and Werb (1996) Curr. Opin. Cell Biol. 8:731-738; Birkedal-Hansen, et al. 30 (1993) Crit. Rev. Oral Biol. Med. 4:197-250; Mignatti and Rifkin (1993) Physiol. Rev. 73:161-195; and Stetler-Stevenson, et al. (1993) Annu. Rev. Cell Biol. 9:541-573. They play crucial roles in tissue remodeling in normal  
35 and pathological processes, including development, repair, and cancer progression. All MMPs identified to date are synthesized as an inactive proenzyme form or zymogen, contain zinc-binding sites, and need proteolytic activation to become functional proteases. According to

their structural features and substrate specificity, four subclasses of MMPs have been established: collagenases have the unique capacity to degrade fibrillar collagens; gelatinases denature basement membranes and denatured collagens; stromelysins degrade many extracellular proteins, including proteoglycans, laminin, and fibronectin; and membrane-type MMPs are supposed to have proteolytic activity on other MMPs, required for their activation. See Matrisian (1992) Bioessays. 14:455-463; Stetler-Stevenson, et al. (1993) Annu. Rev. Cell Biol. 9:541-573; and Woessner (1991) FASEB J. 5:2145-2154. Among the 15 identified MMPs, four distinct members, presenting a transmembrane domain at the C terminus, have been described and are referred to as MT-MMP (Membrane-type matrix metalloproteinase). Sato, et al. (1994) Nature 370:61-65 first identified the MT1-MMP (MMP 14), responsible for the activation of progelatinase A (pro-MMP2) on the tumor cell surface that may trigger tissue invasion by tumor cells. Three additional members of this family, MT2-MMP (MMP 15), MT3-MMP (MMP 16) and MT4-MMP (MMP 17) have been isolated respectively from lung, placenta, and breast carcinoma cDNA libraries. See Puente, et al. (1996) Cancer Res. 56:944-949; Takino, et al. (1995) J. Biol. Chem. 270:23013-23020; and Will and Hinzmann (1995) Eur. J. Biochem. 231:602-608. All MT-MMPs, like the stromelysin-3 MMP (Basset, et al. (1990) Nature 348:699-704), contain a consensus insertion of about ten amino acids (RxK/RR) between the propeptide and the catalytic domain, corresponding to potential cleavage sites by enzymes called furin (Basbaum and Werb (1996) Curr. Opin. Cell Biol. 8:731-8; Sang and Douglas (1996) J. Protein Chem. 15:137-160). This cleavage is necessary to give rise to an activated form of MT-MMPs. The MT-MMPs are located in ternary complexes including a substrate, a tissue inhibitor of MMPs (TIMPs), and an activated MT-MMP, associated with the plasma membrane (Stetler-Stevenson, et al. (1993) Annu. Rev. Cell Biol. 9:541-573). As described for MT1-, MT2-, and MT3-MMPs, MT-MMPs may have a proteolytic activity on other MMPs

like gelatinase A (pro-MMP2) and collagenase-3 (MMP 13) (see Butler, et al. (1997) Eur. J. Biochem. 244:653-657; Knauper, et al. (1996) J. Biol. Chem. 271:17124-17131; Kolkenbrock, et al. (1997) Biol. Chem. 378:71-76; Sato, et al. (1994) Nature 370:61-65; Strongin, et al. (1993) J. Biol. Chem. 268:14033-14039; and Takino, et al. (1995) J. Biol. Chem. 270:23013-23020).

The present invention provides DNA sequences encoding mammalian proteins which exhibit structural properties or motifs characteristic of a protease, more particularly a matrix metalloproteinase. The proteins described herein are designated F06B09. A human F06B09 nucleotide and predicted amino acid sequence are provided in SEQ ID NO: 1 and 2. The signal sequence/cleavage is predicted to be at nucleotide 344. The predicted extracellular domain is expected at about 1-527; the transmembrane segment occurs at about 528-543; the cytoplasmic domain is found at about 544-545.

An alternative sequence of F06B09 is provided in SEQ ID NO: 3 and 4. Notable motifs include: predicted signal sequence until about nucleotide 343; propeptide domain from about 1-66; C switch motif from about 67-73; furin site from about 82-86; catalytic site from about 87-211; zinc binding site from about 212-222, with notable His at 212, 216, and 222; hinge region from about 260-290; hemopexin-like domain from about 291-525; transmembrane segment from about 526-538; and cytoplasmic tail from about 539-542.

Through an effort aiming at the identification of human dendritic cells (DC) specific genes, the cDNA coding for a fifth member of the human Membrane-type Matrix Metalloproteinases (MT-MMP) family has been cloned. The full-length 3691 bp cDNA which was mapped on chromosome 16p13.3, contains an open reading frame of some 1689 bp, encoding a 562 amino acid protein. The predicted protein was most homologous (48% amino acid homology) with the human matrix metalloproteinase MT4-MMP and has the typical features of member of the MMP family,



WO 99/35276

including a prodomain with the activation locus, the zinc binding site, and the hemopexin domain.

The general roles of matrix metalloproteases are described above. The specific interaction of matrix metalloproteinases with other proteins, e.g., furin and progelatinase, are described in Basbaum and Werb (1996) Current Opinion in Cell Biology 8:731-738. Matrix metalloproteases are typically zinc endopeptidases that are required for the degradation of extracellular matrix components during normal embryo development, morphogenesis, and tissue remodeling. Their proteolytic activities are precisely regulated by endogenous tissue inhibitors of metalloproteases (TIMPS). Disruption of this balance results in diseases such as arthritis, atherosclerosis, and tumor growth and metastasis. Nagase (1996) in Hooper (ed.) Zinc Metalloproteinases in Health and Disease Taylor and Francis, London; Coussens and Werb (1996) Chem. Biol. 3:895-904. Therefore, F06B09 gene product could play a role in the migration of the dendritic cells (DC) or in the progression of the dendrites between the stromal cells. The way the MMPs act on the matrix is complex. The MMP is typically produced as an inactive proenzyme that needs to be processed by another protease, most probably furin, since F06B09 contains a site of cleavage for this convertase. This process probably occurs intracellularly (Basbaum and Werb (1996) Current Opinion in Cell Biology 8:731-738), and is likely followed by an interaction of F06B09 with other proteases like progelatinase.

A single 3.7 Kb mRNA transcript of this gene was found to be mainly expressed in CD34<sup>+</sup>-derived human DC and also weakly in in vitro generated granulocytes. No signal was detected in TF1, CHA, Jurkat, MRC5, or U937 cell lines, nor in freshly isolated monocytes, activated T and B cells, and activated peripheral blood lymphocytes (PBLs). Among normal adult human tissues, this mRNA was detected in spleen, lymph node, thymus, appendix, and bone marrow, but no expression was found in fetal tissues. RT-PCR distribution analysis showed a

significant expression of the novel MT-MMP in activated DC and weakly in JY B cell line. Interestingly, it was found that the novel MT-MMP mRNA expression was down-regulated upon DC activation with CD40L. The expression  
5 pattern of this gene, which is predominantly expressed by DC, together with its putative membrane localization, suggest that it could be involved in the degradation of the extracellular matrix during DC migration.

The descriptions below are directed, for exemplary  
10 purposes, to primate embodiments, e.g., human, but are likewise applicable to related embodiments from other, e.g., natural, sources. Other ESTs have been identified from rodent cDNA libraries. These sources should, where appropriate, include various vertebrates, typically warm  
15 blooded animals, e.g., birds and mammals, particularly domestic animals, and primates. The sequences exhibit significant similarity to membrane-type matrix metalloproteases MT-MMP1 to 4. Table 1 shows an  
20 alignment of the family members.

Table 1: Comparison of various MMPs with F06B09. MT4-MMP (SEQ ID NO: 7) is from Genbank 3466295 (EMBL X89576); MT2-MMP (SEQ ID NO: 8) is from Genbank 1418215 (DDBJ D86331; see also Genbank 3646295 (EMBL Z48482); MT1-MMP (SEQ ID NO: 9) is from Genbank 804994 (EMBL X83535; see also Genbank 1495995 (EMBL X90925) and 793763 (DDBJ D26512); and MT3-MMP (SEQ ID NO: 10) is from Genbank 2424979 (DDBJ D85511).

	MT4-MMP	1			0
10	F06B09	1	MRLRLRLALLLLLLLAPPARAPKPSAQDVS LGVDWLTRY		39
	MT2-MMP	1			0
	MT1-MMP	1	MSPAPRPSRCLLLPLLTGTALASLGSAQSSSFSP EAWLQQY		42
	MT3-MMP	1	MILLTFSTGRRLDFVHSGVFFLQTLWLILCATVCGTEQYFNVEVWLQKY		50
15	MT4-MMP	1		MQQFGGLEATGILDEATLALMKTTPR	25
	F06B09	40	GYLPPPHPAQAQLQSPEKLRDAIKVMQRFAGLPETGRMDPGTVATMRKPR		89
	MT2-MMP	1		MKRPR	5
	MT1-MMP	43	GYLPPGDLRTHTRSPQSLSAIAAMQKFYGLQVTGKADATMKAMRRPR		92
20	MT3-MMP	51	GYLPPTDPRMSVLRSAETMQSALAAMQQFYGINMTGKVDRTIDWMKKPR		100
			* . . *		
	MT4-MMP	26	CSLPDLP-VLTQARRR--RQ--APAPTKWNKRNL SWRVRTFPRDSP LGHD		70
	F06B09	90	CSLPDVL-GVAGLVRR--RRRYALSGSVWKKRTLTWRVRSFPQSSQLSQE		136
25	MT2-MMP	6	CGVPDQFGVRVKANLRRRRKRYALTGRKWNHHLTF SIQNYT--EKL GWY		53
	MT1-MMP	93	CGVPDKFGAEIKANVR--RKRYAIQGLKWQHNEITFCIQNYT--PKVGEY		138
	MT3-MMP	101	CGVPDQTRGSSKFHIR--RKRYALTGQKWQHKHITYSIK NVT--PKVGDP		146
			* . . *		
30	MT4-MMP	71	TVRALMYIALKVWSDIAPLNFHEVA---GS-----TADIQIDFSKADHND		112
	F06B09	137	TVRVLMSYALMAWGMESGLTFHEVDSPQGQ-----EPDILIDFARAFHQD		181
	MT2-MMP	54	HSMEAVRRAFRVWEQATPLVFQEVPIEDIRLRROKEADIMVLFASGFHGD		103
	MT1-MMP	139	ATYEAIRKAFRVWESATPLRFREVPYAYIREGHEKQADIMIFFAEGFHGD		188
	MT3-MMP	147	ETRKAIRRAFVDVWQNVPLTFEEVPYSELENGK-RDVDITIIFASGFHGD		195
35			* . . *		
	MT4-MMP	113	GYPFDGPGGTVAHAFPPGHHTAGDTHFDDDEAWTFRSSDAHGMDFLFAVA		162
	F06B09	182	SYPF DGLGGLAHAFPPGEHPISGDTHFDDDEETWTFGSKDGEGLDFLFAVA		231
	MT2-MMP	104	SSPFDGTGGFLAHAYFP GPG-LGGDTHFDADEPWTFSSTDLHGNNLFLVA		152
40	MT1-MMP	189	STPFDGEGGFLAHAYFP GPN-IGGDTHFDSAEPWTVRNLGNDIFLVA		237
	MT3-MMP	196	SSPFDGEGGFLAHAYFP GPG-IGGDTHFDSDEPWT LGNPNHGDNDLFLVA		244
			**** * . . *		
45	MT4-MMP	163	VHEFGHAIGLSHVAAHSIMRPPYQGPVGDPLRYGLPYEDKVRVWQLYGV		212
	F06B09	232	VHEFGHALGLGHSSAPNSIMRPFYQGPVGD PKYRLSQDDRDGLQQLYG-		280
	MT2-MMP	153	VHELGHALGLEHSSNPNAIMAPFYQWKDNDN--FKLPEDDLRGIQQLYGT		200
	MT1-MMP	238	VHELGHALGLEHSSDPSAIMAPFYQWMDTEN--FVLPDDDRRGIQQLYGG		285
	MT3-MMP	245	VHELGHALGLEHSNDPTAIMAPFYQYMETDN--FKLPNDLQGIQKIYGP		292
			*** * . . *		
50	MT4-MMP	213	RESVSPTAQ--PEEPPLLP-----EP-----PDNRSSA		238
	F06B09	281	KAPQTPYDK--PTRKPLAP-----PPQ-----PPASPTH		307
	MT2-MMP	201	PDGQPQPTQPLPTVTPRRPG-----RPDHRPPRPQP PPPPGKPERPPKP		245
	MT1-MMP	286	ESG-----FPTKMPPQP-----RTTSRP-----SVPDKPKNP		312
55	MT3-MMP	293	PKI PPPTRPLPTVPPHRSIPPADPRKNDRP-----KPPRPPTG		331
			* . . *		

5	MT4-MMP	239	PPR-----KDVPHRCSTHFDVAQIRGEAFFFKGKYFWRLTRDRH	278
	F06B09	308	SPS-----FPIPDRCENFDAIANIRGETFFFKGPWFWRQLQPSGQ	347
	MT2-MMP	246	GPPVQPRATERPDQYGNICDGDFTVAMLRGEMFVFKGRWFWVRVHRNR-	294
	MT1-MMP	313	-----TYGPNICDGNFDTVAMLRGEMFVFKERWFWVRVNNQ-	348
	MT3-MMP	332	RPS-----YPGAKPNICDGNFNTLAILRREMFMFKDQWFWVRVNNR-	372
* * * . . * * * * * *				
10	MT4-MMP	279	LVS LQPAQMHRFWRGLPLHLDSVDAVYERTSDHKIVFFKGDYRWVFKDNN	328
	F06B09	348	LVSPRPARLHRFWEGFLPAQVRVVQAAYARHRDGRILLFSGPQFWVFQDR-	396
	MT2-MMP	295	VLDNYPMPIGHFWRGLPGDIS---AAYERQ-DGRFVFFKGDYRWLFREAN	340
	MT1-MMP	349	VMDGYPMPIGQFWRGLPASIN---TAYERK-DGKFVFFKGDKHWVFDEAS	394
	MT3-MMP	373	VMDGYPMQITYFWRGLPPSID---AVYENS-DGNFVFFKGNKYWVFKDTT	418
.. * . * * * * . . * * . * * . * . *				
15	MT4-MMP	329	VEEGYPRPVSDFSLP--PGG-IDAAFSAWHNDRTYFFKDQLYWRYDDHTR	375
	F06B09	397	QLEGGARPLTELGLP--PGEEDAVFSWPQNGKTYLVRGRQYWRYDEAAA	444
	MT2-MMP	341	LEPGYPQPLTSYGL-GIPYDRIDTAIWWEPTGHTFFFQEDRYWRFNEETQ	389
	MT1-MMP	395	LEPGYPKHikelGR-GLPTDKIDAALFWMPNGKTYFFRGNKYYRFNEELR	443
	MT3-MMP	419	LQPGYPHDLITLGS-GIPPHGIDSaiWWEDVGKTYFFKGDYRWRYSEEMK	467
* . . . * . * . * . * . * . *				
20	MT4-MMP	376	HMDPGYPAQSPLWRGVPSTLDDAMRWS-DGASYFFRGQYWKVLDGELEV	424
	F06B09	445	RPDPGYPRDLSLWEGAPSPDDVTVSN-AGDTYFFKGAHYWRFPKNSIKT	493
	MT2-MMP	390	RGDPGYPKPISVWQGI PASPKGAFLSNDAAITYFYKGTKYWKFDNERLRM	439
	MT1-MMP	444	AVDSEYPKNIKVWEGIPESPRGSFMSGDEVFTYFYKGNKYWKFNQKLKV	493
	MT3-MMP	468	TMDPGYPKPITVWKGIPESPOGAFVHKENGFTYFYKGKEYWKFNQILKV	517
* ** . * * * . * . * * *				
25	MT4-MMP	425	APGYPQSTARDWLVCGDSQADGSVAAGV-----DAAEGPRAPPGQHDQ	467
	F06B09	494	EPDAPQPMGPNWLDPCAP-----SSGPRAP---RP	520
	MT2-MMP	440	EPGYPKSILRDFMGCEHVEPGPRWPDVARPPFNPHGGAEPGADSAGEDV	489
	MT1-MMP	494	EPGYPKSALRDWMGCPSGGRPDE-----GTEEETEVIIEV	529
	MT3-MMP	518	EPGYPRSILKDFMGCDGPTDRVKEG-----HSPDDVDIVIKL	555
* * . . *				
30	MT4-MMP	468	SRS-----EDGYEVCSTSGASSPPGAPGPLVAAT	497
	F06B09	521	PKA-----TPVSETCDCQCELN---QAAGRWPAPI	547
	MT2-MMP	490	GDGDGDFGAGVKNDRGSRVVVQMEEVARTVNVVMVLVPLLLLLLCVLGLTY	539
	MT1-MMP	530	D-----EEGGGAVSAAAVLPLVLLLLLVLAUGL	557
	MT3-MMP	556	DN-----TASTVKAIAIVIPCILALCLLVLY	582
40	MT4-MMP	498	MLLLLP-PLSPGALWTAQAALTL	519
	F06B09	548	PLLLLP-LLVGGVASR	562
	MT2-MMP	540	ALVQMQRKGAPRVLLYCKRSLQEWV	564
	MT1-MMP	558	AVFFFRRHGTPRRLLYCQRSLLDKV	582
	MT3-MMP	583	TVFQFKRKGTPRHILYCKRSMQEWV	607
45	MT4-MMP	498	MLLLLP-PLSPGALWTAQAALTL	519
	F06B09	548	PLLLLP-LLVGGVASR	562
	MT2-MMP	540	ALVQMQRKGAPRVLLYCKRSLQEWV	564
	MT1-MMP	558	AVFFFRRHGTPRRLLYCQRSLLDKV	582
	MT3-MMP	583	TVFQFKRKGTPRHILYCKRSMQEWV	607
50	MT4-MMP	498	MLLLLP-PLSPGALWTAQAALTL	519
	F06B09	548	PLLLLP-LLVGGVASR	562
	MT2-MMP	540	ALVQMQRKGAPRVLLYCKRSLQEWV	564
	MT1-MMP	558	AVFFFRRHGTPRRLLYCQRSLLDKV	582
	MT3-MMP	583	TVFQFKRKGTPRHILYCKRSMQEWV	607

In the search for DC specific genes, a novel matrix-metalloproteinase homologue (MMP) from the Membrane-type Matrix Metalloproteinases (MT-MMP) family subclass was identified. Of interest, this novel gene designated F06B09 is predominantly expressed by both CD34<sup>+</sup>- and monocyte-derived DC and is down-regulated after DC maturation by CD40L L cells. The deduced protein sequence of F06B09 is clearly a member of the MMP family, characterized by the presence of a prodomain with the activation locus containing the essential cysteine residue, a catalytic domain including the zinc-binding site with the consensus sequence HExGHxxxxxH and an hemopexin-like domain (Birkedal-Hansen (1995) Curr. Opin. Cell Biol. 7:728-735; Shapiro (1998) Curr. Opin. Cell Biol. 10:602-608). MMPs belonging to the metzincin (or Clan) superfamily, can be classified into at least four subfamilies of closely related members: collagenases, stromelysins, gelatinases, and MT-MMP, although there are some MMPs like the macrophage metalloelastase (Belaouaj, et al. (1995) J. Biol. Chem. 270:14568-14575) and the stromelysin 3 (Basset, et al. (1990) Nature 348:699-704) that do not belong to these subclasses. According to its structural characteristic and its high level of homology with MT4-MMP (Puente, et al. (1996) Cancer Res. 56:944-949), F06B09 represents the fifth member of the MT-MMP subclass. All MT-MMPs present a putative transmembrane domain in the C-terminal portion and a characteristic insertion between the propeptide and the catalytic domain containing the consensus amino acid sequence RxK/RR (Basbaum and Werb (1996) Curr. Opin. Cell Biol. 8:731-738). In order to activate MT-MMPs, this site is cleaved by enzymes called furin. Like other MMPs or matrixins, the novel MT-MMP contains noncatalytic domains, in addition to the protease domain, which are likely involved in interactions with substrates or other proteins. MT-MMPs have the ability to cleave substrates, e.g., other MMPs. Effectively, MT1-, MT2- and MT3-MMP can activate proMMP2 (progelatinase A) into MMP2 on the cell surface (Butler, et al. (1997) Eur. J. Biochem.

244:653-657; Knauper, et al. (1996) J. Biol. Chem. 271:17124-17131; Kolkenbrock, et al. (1997) Biol. Chem. 378:71-76; Sato, et al. (1994) Nature 370:61-65; Strongin, et al. (1993) J. Biol. Chem. 268:14033-14039; and Takino, et al. (1995) J. Biol. Chem. 270:23013-23020), which as an active form degrades type IV collagen, the major component of basement membranes (Wilhelm, et al. (1989) J. Biol. Chem. 264:17213-17221). Thus, active MMP-2 (gelatinase A) plays a key role in the invasion of migrating cells into tissues. Several reports demonstrate that an overexpression of MMPs, especially MMP-2 and MMP-9, is associated with the invasive behavior of tumor cells (Shapiro (1998) Curr. Opin. Cell Biol. 10:602-608; Stetler-Stevenson, et al. (1993) Annu. Rev. Cell Biol. 9:541-573). Concerning the MT4-MMP, the most homologous gene to F06B09, it is uncertain whether this MT-MMP can activate proMMP-2. Phylogenetic analysis shows that among the MT-MMPs, MT4-MMP and F06B09 are distinguished from others and form a group of closely related proteins. It is speculated that these two MT-MMP share similar biological activities. It has been shown that MT-MMPs are overexpressed in cancers; in particular, high levels of MT1-MMP are associated with invasiveness of cervical cancer cells (Gilles, et al. (1996) Int. J. Cancer. 65:209-213), breast, colon, neck and lung carcinomas (Okada, et al. (1995) Proc. Natl. Acad. Sci. USA. 92:2730-2734; Sato, et al. (1994) Nature 370:61-65; Ueno, et al. (1997) Cancer Res. 57:2055-2060) and gastric cancers (Mori, et al. (1997) Int. J. Cancer. 74:316-321). Furthermore, MT4-MMP was isolated from a breast carcinoma (Puente, et al. (1996) Cancer Res. 56:944-949). Interestingly, the novel MT-MMP identified in the present report constitutes the first MT-MMP isolated from a DC library. It will be interesting to further study F06B09 expression in pathological tissues.

MT-MMP participate not only to the control of cell migration and tissue remodeling through their involvement in degradation of extracellular matrix and basement

membranes, but also in the processing of pro-enzymes and pro-cytokines.

Of note, the genes for human MT1-MMP, 2, and 3 have been localized on three different chromosomes by in situ hybridization (Mattei, et al. (1997) Genomics. 40:168-169; Mignon, et al. (1995) Genomics. 28:360-361), and while the novel MT-MMP gene is on the same chromosome as MT2-MMP, both genes are on different loci: MT2-MMP is on chromosome 16q12 (Mattei, et al. (1997) Genomics. 40:168-169; Yasumitsu, et al. (1997) DNA Res. 4:77-79), and the novel MT-MMP is on chromosome 16p13.3.

The predominant expression of F06B09 in immature DC, and its putative membrane localization, suggests a role for this MT-MMP in degradation of extracellular matrix during DC migration. Furthermore, like MT1-MMP, which may trigger invasion by tumor cells by activating progelatinase A on tumor cell surface, this novel MT-MMP could be involved in cancer invasion.

The proteins of this invention are defined in part by their sequences, and by their physicochemical and biological properties. The biological properties of the human proteases described herein, e.g., human F06B09, are defined by their amino acid sequences, and mature sizes. They also should share certain biological enzymatic properties of their respective proteins.

The human protease F06B09 translation product exhibits structural motifs of a member of the matrix metalloproteinase family of proteases, more specifically to a family of matrix degrading proteinases. These proteins, in the latent form, typically possess a prodomain form which masks the catalytic site, which chelates a zinc ion. See Vallee and Auld (1990) Biochemistry 29:5647-5659. The processed mature protein is typically a potent cell-matrix degrading enzyme. See, e.g., Birkedal-Hansen (1990) Proc. Nat'l Acad. Sci. USA 87:5578-5582. The enzyme may remain attached to the cell membrane after activation, and many of these proteases may localize to the leading cellular processes when the cell migrates. This may suggest other protein-protein

interactions, e.g., with domains which specifically localize the enzymes by cytoskeletal or other mechanisms.

The pro-enzyme activation (furin) site would correspond to the Arg stretch from 86-90 of SEQ ID NO: 2; or 212-222 of SEQ ID NO: 4. It is likely that the activating enzyme will be one of the furin/PACE proteases, which are essentially ubiquitously expressed. By analogy to the MT1-MMP, these proteases activate other proteases, e.g., gelatinase A, collagenases, and others, that assist in the degradation of the extracellular matrix. See, e.g., Basbaum and Werb (1996) Current Opinion in Cell Biology 8:731-738.

F06B09 contains the zinc binding peptide consensus at about residues 212-222 of SEQ ID NO: 4, with characteristic His residues at 212, 216, and 222. There is also a hemopexin-like domain at about residues 291-525 and a matrixin-like domain corresponding to about residues 1-211. Natural substrates for the proteinase may be identified using standard methods. Substrate sequence specificity may be determined, and search for such sequences in databases may identify specific candidates for physiological substrates.

One of skill will readily recognize that some sequence variations may be tolerated, e.g., conservative substitutions or positions remote from the critical helical structures and remote from the identified or consensus critical active site regions, without altering significantly the biological activity of each respective molecule.

F06B09 proteins are present in specific cell types, e.g., dendritic cells, and the interaction of the protease with a substrate will be important for mediating various aspects of cellular physiology or development. The cellular types which express messages encoding F06B09 suggest that signals important in cell differentiation and development are mediated by them. See, e.g., Gilbert (1991) Developmental Biology (3d ed.) Sinauer Associates, Sunderland, MA; Browder, et al. (1991) Developmental Biology (3d ed.) Saunders, Philadelphia, PA.; Russo, et



al. (1992) Development: The Molecular Genetic Approach  
Springer-Verlag, New York, N.Y.; and Wilkins (1993)  
Genetic Analysis of Animal Development (2d ed.) Wiley-  
Liss, New York, N.Y. In particular, the proteases may be  
5 necessary for the conversion of pro-proteins to proteins,  
e.g., cytokine or protein precursors to mature forms, or  
for proper immunological function, e.g., antigen  
processing and presentation. Alternatively, the  
proteases may be important in dendritic cell trafficking,  
10 e.g., to traverse through extracellular matrix or  
vascular surfaces.

## II. Definitions

The term "binding composition" refers to molecules  
15 that bind with specificity to F06B09, e.g., in an  
antibody-antigen interaction. However, other compounds,  
e.g., complex associated proteins, may also specifically  
associate with F06B09 to the exclusion of other  
molecules. Typically, the association will be in a  
20 natural physiologically relevant protein-protein  
interaction, either covalent or non-covalent, and may  
include members of a multiprotein complex, including  
carrier compounds or dimerization partners. The molecule  
may be a polymer, or chemical reagent. A functional  
25 analog may be a protease with structural modifications,  
or may be a wholly unrelated molecule, e.g., which has a  
molecular shape which interacts with the appropriate  
substrate cleavage determinants.

The term "binding agent:F06B09 protein complex", as  
30 used herein, refers to a complex of a binding agent and  
an F06B09 protein that is formed by specific binding of  
the binding agent to the F06B09 protein. Specific  
binding of the binding agent means that the binding agent  
has a specific binding site that recognizes a site on the  
35 F06B09 protein, typically in the native conformation, but  
possibly in a denatured conformation, e.g., a Western  
blot. For example, antibodies raised to an F06B09  
protein and recognizing an epitope on the F06B09 protein  
are capable of forming a binding agent:F06B09 protein

complex by specific binding. Typically, the formation of a binding agent:F06B09 protein complex allows the measurement of F06B09 protein in a biological sample, e.g., a mixture with other proteins and biologics. The  
5 term "antibody:F06B09 protein complex" refers to an embodiment in which the binding agent is an antibody. The antibody may be monoclonal, polyclonal, or a binding fragment of an antibody, e.g., an Fab, F(ab)2, or Fv fragment. The antibody will preferably be a polyclonal  
10 antibody for cross-reactivity determinations.

"Homologous" nucleic acid sequences, when compared, exhibit significant similarity or identity. The standards for homology in nucleic acids are either measures for homology generally used in the art by  
15 sequence comparison and/or phylogenetic relationship, or based upon hybridization conditions. Hybridization conditions are described in greater detail below.

An "isolated" nucleic acid is a nucleic acid, e.g., an RNA, DNA, or a mixed polymer, which is substantially  
20 separated from other biologic components which naturally accompany a native sequence, e.g., proteins and flanking genomic sequences from the originating species. The term embraces a nucleic acid sequence which has been removed from its naturally occurring environment, and includes  
25 recombinant or cloned DNA isolates and chemically synthesized analogs, or analogs biologically synthesized by heterologous systems. A substantially pure molecule includes isolated forms of the molecule. An isolated nucleic acid will usually contain homogeneous nucleic  
30 acid molecules, but will, in some embodiments, contain nucleic acids with minor sequence heterogeneity. This heterogeneity is typically found at the polymer ends or portions not critical to a desired biological function or activity.

35 As used herein, the term "F06B09" protein shall encompass, when used in a protein context, a protein having amino acid sequences, particularly from the protein motif portions, shown in SEQ ID NO: 2 or 4, respectively. In many contexts, a significant fragment

of such a protein will be functionally equivalent. The invention also embraces a polypeptide which exhibits similar structure to human F06B09 protein, e.g., which interacts with F06B09 specific binding components. These  
5 binding components, e.g., antibodies, typically bind to F06B09 protein with high affinity, e.g., at least about 100 nM, usually better than about 30 nM, preferably better than about 10 nM, and more preferably at better than about 3 nM.

10 The term "polypeptide" or "protein" as used herein includes a significant fragment or segment of protease motif portion of F06B09 protein, and encompasses a stretch of amino acid residues of at least about 8 amino acids, generally at least about 10 amino acids, more  
15 generally at least about 12 amino acids, often at least about 14 amino acids, more often at least about 16 amino acids, typically at least about 18 amino acids, more typically at least about 20 amino acids, usually at least about 22 amino acids, more usually at least about 24  
20 amino acids, preferably at least about 26 amino acids, more preferably at least about 28 amino acids, and, in particularly preferred embodiments, at least about 30 or more amino acids, e.g., 35, 40, 45, 50, 60, 70, 80, 100, etc. Preferred ends of such polypeptides will correspond  
25 to a motif or boundary described above, e.g., in Table 1. Preferably, a polypeptide will contain a plurality of distinct, e.g., discrete or nonoverlapping, segments of the specified length. Typically, the plurality will be at least two, more usually at least three, and preferably  
30 5, 7, or even more. While the length minima are provided, longer lengths, of various sizes, may be appropriate, e.g., one of length 7, and two of length 12.

A "recombinant" nucleic acid is defined either by its method of production or its structure. In reference  
35 to its method of production, e.g., a product made by a process, the process is use of recombinant nucleic acid techniques, e.g., involving human intervention in the nucleotide sequence, typically selection or production. Alternatively, it can be a nucleic acid made by

generating a sequence comprising fusion of two fragments which are not naturally contiguous to each other, but is meant to exclude products of nature, e.g., naturally occurring mutants. Thus, for example, products made by transforming cells with any non-naturally occurring vector is encompassed, as are nucleic acids comprising sequence derived using any synthetic oligonucleotide process. Such is often done to replace a codon with a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a sequence recognition site. Alternatively, it is performed to join together nucleic acid segments of desired functions to generate a single genetic entity comprising a desired combination of functions not found in the commonly available natural forms. Restriction enzyme recognition sites are often the target of such artificial manipulations, but other site specific targets, e.g., promoters, DNA replication sites, regulation sequences, control sequences, or other useful features may be incorporated by design. A similar concept is intended for a recombinant, e.g., fusion, polypeptide. Specifically included are synthetic nucleic acids which, by genetic code redundancy, encode polypeptides similar to fragments of these antigens, and fusions of sequences from various different species variants.

"Solubility" is reflected by sedimentation measured in Svedberg units, which are a measure of the sedimentation velocity of a molecule under particular conditions. The determination of the sedimentation velocity was classically performed in an analytical ultracentrifuge, but is typically now performed in a standard ultracentrifuge. See, Freifelder (1982) Physical Biochemistry (2d ed.) W.H. Freeman & Co., San Francisco, CA; and Cantor and Schimmel (1980) Biophysical Chemistry parts 1-3, W.H. Freeman & Co., San Francisco, CA. As a crude determination, a sample containing a putatively soluble polypeptide is spun in a standard full sized ultracentrifuge at about 50K rpm for about 10 minutes, and soluble molecules will remain in the

supernatant. A soluble particle or polypeptide will typically be less than about 30S, more typically less than about 15S, usually less than about 10S, more usually less than about 6S, and, in particular embodiments, preferably less than about 4S, and more preferably less than about 3S. Solubility of a polypeptide or fragment depends upon the environment and the polypeptide. Many parameters affect polypeptide solubility, including temperature, electrolyte environment, size and molecular characteristics of the polypeptide, and nature of the solvent. Typically, the temperature at which the polypeptide is used ranges from about 4° C to about 65° C. Usually the temperature at use is greater than about 18° C and more usually greater than about 22° C. For diagnostic purposes, the temperature will usually be about room temperature or warmer, but less than the denaturation temperature of components in the assay. For therapeutic purposes, the temperature will usually be body temperature, typically about 37° C for humans, though under certain situations the temperature may be raised or lowered in situ or in vitro.

The size and structure of the polypeptide should generally be in a substantially stable state, and usually not in a denatured state. The polypeptide may be associated with other polypeptides in a quaternary structure, e.g., to confer solubility, or associated with lipids or detergents in a manner which approximates natural lipid bilayer interactions.

The solvent will usually be a biologically compatible buffer, of a type used for preservation of biological activities, and will usually approximate a physiological solvent. Usually the solvent will have a neutral pH, typically between about 5 and 10, and preferably about 7.5. On some occasions, a detergent will be added, typically a mild non-denaturing one, e.g., CHS (cholesteryl hemisuccinate) or CHAPS (3-[3-cholamidopropyl]-dimethylammonio]-1-propane sulfonate), or a low enough concentration as to avoid significant

disruption of structural or physiological properties of the protein.

"Substantially pure" in a protein context typically means that the protein is isolated from other  
5 contaminating proteins, nucleic acids, and other biologicals derived from the original source organism. Purity, or "isolation" may be assayed by standard methods, and will ordinarily be at least about 50% pure, more ordinarily at least about 60% pure, generally at  
10 least about 70% pure, more generally at least about 80% pure, often at least about 85% pure, more often at least about 90% pure, preferably at least about 95% pure, more preferably at least about 98% pure, and in most preferred  
15 embodiments, at least 99% pure. Similar concepts apply, e.g., to antibodies or nucleic acids.

"Substantial similarity" in the nucleic acid sequence comparison context means either that the segments, or their complementary strands, when compared, are identical when optimally aligned, with appropriate  
20 nucleotide insertions or deletions, in at least about 50% of the nucleotides, generally at least about 56%, more generally at least about 59%, ordinarily at least about 62%, more ordinarily at least about 65%, often at least about 68%, more often at least about 71%, typically at  
25 least about 74%, more typically at least about 77%, usually at least about 80%, more usually at least about 85%, preferably at least about 90%, more preferably at least about 95 to 98% or more, and in particular  
30 embodiments, as high at about 99% or more of the nucleotides. Alternatively, substantial similarity exists when the segments will hybridize under selective hybridization conditions, to a strand, or its complement, typically using a sequence derived from SEQ ID NO: 1 or  
3. Typically, selective hybridization will occur when  
35 there is at least about 55% similarity over a stretch of at least about 30 nucleotides, preferably at least about 65% over a stretch of at least about 25 nucleotides, more preferably at least about 75%, and most preferably at least about 90% over about 20 nucleotides. See Kanehisa

(1984) Nucl. Acids Res. 12:203-213. The length of similarity comparison, as described, may be over longer stretches, and in certain embodiments will be over a stretch of at least about 17 nucleotides, usually at least about 20 nucleotides, more usually at least about 24 nucleotides, typically at least about 28 nucleotides, more typically at least about 40 nucleotides, preferably at least about 50 nucleotides, and more preferably at least about 75 to 100 or more nucleotides, e.g., 150, 200, etc. Various combinations of plurality of such segments will also be made.

For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

Optical alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith and Waterman (1981) Adv. Appl. Math. 2:482, by the homology alignment algorithm of Needleman and Wunsch (1970) J. Mol. Biol. 48:443, by the search for similarity method of Pearson and Lipman (1988) Proc. Nat'l Acad. Sci. USA 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by visual inspection (see generally Ausubel et al., supra).

One example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a tree or dendrogram showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of

Feng and Doolittle (1987) J. Mol. Evol. 35:351-360. The method used is similar to the method described by Higgins and Sharp (1989) CABIOS 5:151-153. The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison and by designating the program parameters. For example, a reference sequence can be compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps.

Another example of algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described Altschul, et al. (1990) J. Mol. Biol. 215:403-410. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul, et al., supra). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased.



Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a wordlength (W) of 11, the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) Proc. Nat'l Acad. Sci. USA 89:10915) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin and Altschul (1993) Proc. Nat'l Acad. Sci. USA 90:5873-5787). One measure of similarity provided by the BLAST algorithm is the smallest sum probability ( $P(N)$ ), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

A further indication that two nucleic acid sequences of polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second polypeptide, for example, where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules hybridize to each other under stringent conditions, as described below.

"Stringent conditions", in referring to homology or substantial similarity in the hybridization context, will be stringent combined conditions of salt, temperature, organic solvents, and other parameters, typically those controlled in hybridization reactions. The combination of parameters is generally more important than the measure of any single parameter. See, e.g., Wetmur and Davidson (1968) J. Mol. Biol. 31:349-370. A nucleic acid probe which binds to a target nucleic acid under stringent conditions is specific for said target nucleic acid. Such a probe is typically more than 11 nucleotides in length, and is sufficiently identical or complementary to a target nucleic acid over the region specified by the sequence of the probe to bind the target under stringent hybridization conditions. Hybridization under stringent conditions should give a background of at least 2-fold over background, preferably at least 3-5 or more.

F06B09 proteins from other mammalian species can be cloned and isolated by cross-species hybridization of closely related species. See, e.g., below. Similarity may be relatively low between distantly related species, and thus hybridization of relatively closely related species is advisable. Alternatively, preparation of an antibody preparation which exhibits less species specificity may be useful in expression cloning approaches.

The phrase "specifically binds to an antibody" or "specifically immunoreactive with", when referring to a protein or peptide, refers to a binding reaction which is determinative of the presence of the protein in the presence of a heterogeneous population of proteins and other biological components. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein and do not significantly bind other proteins present in the sample. Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. For example, antibodies raised to the human F06B09 protein immunogen with the amino acid sequence

depicted in SEQ ID NO: 2 or 4 can be selected by immunoaffinity or similar methods to obtain antibodies specifically immunoreactive with F06B09 proteins and not with other proteins.

5

### III. Nucleic Acids

F06B09 proteins are exemplary of larger classes of structurally and functionally related proteins. The F06B09 proteins will typically serve to cleave or process various proteins produced or processed by various cell types, e.g., for antigen presentation. The preferred embodiments, as disclosed, will be useful in standard procedures to isolate genes from different individuals or other species, e.g., warm blooded animals, such as birds and mammals. Cross hybridization will allow isolation of related genes encoding proteins from individuals, strains, or species. A number of different approaches are available to successfully isolate a suitable nucleic acid clone based upon the information provided herein. Southern blot hybridization studies can qualitatively determine the presence of homologous genes in human, monkey, rat, dog, cow, and rabbit genomes under specific hybridization conditions.

Complementary sequences will also be used as probes or primers. Based upon identification of the likely amino terminus, other peptides should be particularly useful, e.g., coupled with anchored vector or poly-A complementary PCR techniques or with complementary DNA of other peptides.

Techniques for nucleic acid manipulation of genes encoding F06B09 proteins, such as subcloning nucleic acid sequences encoding polypeptides into expression vectors, labeling probes, DNA hybridization, and the like are described generally in Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (2nd ed.) Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor Press, NY, which is incorporated herein by reference. This manual is hereinafter referred to as "Sambrook, et al."

There are various methods of isolating DNA sequences encoding F06B09 proteins. For example, DNA is isolated from a genomic or cDNA library using labeled oligonucleotide probes having sequences identical or complementary to the sequences disclosed herein. Full-length probes may be used, or oligonucleotide probes may be generated by comparison of the sequences disclosed. Such probes can be used directly in hybridization assays to isolate DNA encoding F06B09 proteins, or probes can be designed for use in amplification techniques such as PCR, for the isolation of DNA encoding F06B09 proteins.

To prepare a cDNA library, mRNA is isolated from cells, preferably which express high levels of an F06B09 protein. cDNA is prepared from the mRNA and ligated, e.g., into a recombinant vector. The vector is transfected into a recombinant host for propagation, screening, and cloning. Methods for making and screening cDNA libraries are well known. See Gubler and Hoffman (1983) Gene 25:263-269 and Sambrook, et al.

For a genomic library, the DNA can be extracted from tissue, and often either mechanically sheared or enzymatically digested to yield fragments of about 12-20 kb. The fragments are then separated by gradient centrifugation and cloned in bacteriophage lambda vectors. These vectors and phage are packaged in vitro, as described in Sambrook, et al. Recombinant phage are analyzed by plaque hybridization as described in Benton and Davis (1977) Science 196:180-182. Colony hybridization is carried out as generally described in, e.g., Grunstein, et al. (1975) Proc. Natl. Acad. Sci. USA. 72:3961-3965.

DNA encoding an F06B09 protein can be identified in either cDNA or genomic libraries by its ability to hybridize with the nucleic acid probes described herein, e.g., in colony or plaque hybridization assays. The corresponding DNA regions are isolated, e.g., by standard methods familiar to those of skill in the art. See, e.g., Sambrook, et al.

Various methods of amplifying target sequences, such as the polymerase chain reaction, can also be used to prepare DNA encoding F06B09 proteins. Polymerase chain reaction (PCR) technology may be used to amplify such  
5 nucleic acid sequences directly from mRNA, from cDNA, and/or from genomic libraries or cDNA libraries. The isolated sequences encoding F06B09 proteins may also be used as templates for PCR amplification.

Typically, in PCR techniques, oligonucleotide  
10 primers complementary to two flanking regions in the DNA region to be amplified are synthesized. The polymerase chain reaction is then carried out using the two primers. See Innis, et al. (eds. 1990) PCR Protocols: A Guide to Methods and Applications Academic Press, San Diego, CA.  
15 Primers can be selected to amplify the entire regions encoding a full-length human F06B09 protein or to amplify smaller DNA segments, as desired. Once such regions are PCR-amplified, they can be sequenced and oligonucleotide probes can be prepared from sequence obtained using  
20 standard techniques. These probes can then be used to isolate DNA's encoding F06B09 proteins.

Oligonucleotides for use as probes are usually chemically synthesized according to the solid phase phosphoramidite triester method first described by  
25 Beaucage and Carruthers (1983) Tetrahedron Lett. 22(20):1859-1862, or using an automated synthesizer, as described in Needham-VanDevanter, et al. (1984) Nucleic Acids Res. 12:6159-6168. Purification of oligonucleotides is performed e.g., by native acrylamide  
30 gel electrophoresis or by anion-exchange HPLC as described in Pearson and Regnier (1983) J. Chrom. 255:137-149. The sequence of the synthetic oligonucleotide can be verified using, e.g., the chemical degradation method of Maxam, A.M. and Gilbert, W. in  
35 Grossman and Moldave (eds. 1980) Methods in Enzymology 65:499-560, Academic Press, New York.

An isolated nucleic acid encoding a human F06B09 protein was identified. The nucleotide sequence,

corresponding open reading frames, and mature peptides are provided in Table 1 or SEQ ID NO: 1 or 3.

This invention provides isolated DNA or fragments to encode an F06B09 protein or specific fragment thereof.

- 5 In addition, this invention provides isolated or recombinant DNA which encodes a protein or polypeptide, and which is capable of hybridizing under appropriate conditions, e.g., high stringency, with the DNA sequences described herein. Said biologically active protein or
- 10 polypeptide can be a functional protease segment, or fragment, and have an amino acid sequence as disclosed in SEQ ID NO: 2 or 4. Preferred embodiments will be full length natural sequences, from isolates, or proteolytic fragments thereof. Further, this invention contemplates
- 15 the use of isolated or recombinant DNA, or fragments thereof, which encode proteins which exhibit high measures of identity to an F06B09 protein, or which were isolated, e.g., using cDNA encoding an F06B09 protease polypeptide as a probe. The isolated DNA can have the
- 20 respective regulatory sequences in the 5' and 3' flanks, e.g., promoters, enhancers, poly-A addition signals, and others.

#### IV. Making human F06B09 proteins

- 25 DNAs which encode an F06B09 protein, or fragments thereof, can be obtained by chemical synthesis, screening cDNA libraries, or by screening genomic libraries prepared from a wide variety of cell lines or tissue samples.

- 30 These DNAs can be expressed in a wide variety of host cells for the synthesis of a full-length protein or fragments which can in turn, e.g., be used to generate polyclonal or monoclonal antibodies; for binding studies; for construction and expression of modified molecules;
- 35 and for structure/function studies. Each of F06B09, or their fragments, can be expressed in host cells that are transformed or transfected with appropriate expression vectors. These molecules can be substantially purified to be free of protein or cellular contaminants, other

than those derived from the recombinant host, and therefore are particularly useful in pharmaceutical compositions when combined with a pharmaceutically acceptable carrier and/or diluent. The antigen, e.g., F06B09, or portions thereof, may be expressed as fusions with other proteins or possessing an epitope tag.

Expression vectors are typically self-replicating DNA or RNA constructs containing the desired antigen gene or its fragments, usually operably linked to appropriate genetic control elements that are recognized in a suitable host cell. The specific type of control elements necessary to effect expression will depend upon the eventual host cell used. Generally, the genetic control elements can include a prokaryotic promoter system or a eukaryotic promoter expression control system, and typically include a transcriptional promoter, an optional operator to control the onset of transcription, transcription enhancers to elevate the level of mRNA expression, a sequence that encodes a suitable ribosome binding site, and sequences that terminate transcription and translation. Expression vectors also usually contain an origin of replication that allows the vector to replicate independently from the host cell.

The vectors of this invention contain DNAs which encode an F06B09 protein, or a significant fragment thereof, typically encoding, e.g., a biologically active polypeptide, or protein. The DNA can be under the control of a viral promoter and can encode a selection marker. This invention further contemplates use of such expression vectors which are capable of expressing eukaryotic cDNA coding for an F06B09 protein in a prokaryotic or eukaryotic host, where the vector is compatible with the host and where the eukaryotic cDNA coding for the protein is inserted into the vector such that growth of the host containing the vector expresses the cDNA in question. Usually, expression vectors are designed for stable replication in their host cells or for amplification to greatly increase the total number of

copies of the desirable gene per cell. It is not always necessary to require that an expression vector replicate in a host cell, e.g., it is possible to effect transient expression of the protein or its fragments in various  
5 hosts using vectors that do not contain a replication origin that is recognized by the host cell. It is also possible to use vectors that cause integration of an F06B09 protein gene or its fragments into the host DNA by recombination, or to integrate a promoter which controls  
10 expression of an endogenous gene.

Vectors, as used herein, contemplate plasmids, viruses, bacteriophage, integratable DNA fragments, and other vehicles which enable the integration of DNA fragments into the genome of the host. Expression  
15 vectors are specialized vectors which contain genetic control elements that effect expression of operably linked genes. Plasmids are the most commonly used form of vector, but many other forms of vectors which serve an equivalent function are suitable for use herein. See,  
20 e.g., Pouwels, et al. (1985 and Supplements) Cloning Vectors: A Laboratory Manual Elsevier, N.Y.; and Rodriguez, et al. (eds. 1988) Vectors: A Survey of Molecular Cloning Vectors and Their Uses Butterworth, Boston, MA.

Suitable host cells include prokaryotes, lower eukaryotes, and higher eukaryotes. Prokaryotes include both gram negative and gram positive organisms, e.g., E. coli and B. subtilis. Lower eukaryotes include yeasts, e.g., S. cerevisiae and Pichia, and species of the genus  
30 Dictyostelium. Higher eukaryotes include established tissue culture cell lines from animal cells, both of non-mammalian origin, e.g., insect cells, and birds, and of mammalian origin, e.g., human, primates, and rodents.

Prokaryotic host-vector systems include a wide  
35 variety of vectors for many different species. As used herein, E. coli and its vectors will be used generically to include equivalent vectors used in other prokaryotes. A representative vector for amplifying DNA is pBR322 or its derivatives. Vectors that can be used to express



F06B09 proteins or fragments thereof include, but are not limited to, such vectors as those containing the lac promoter (pUC-series); trp promoter (pBR322-trp); Ipp promoter (the pIN-series); lambda-pP or pR promoters (pOTS); or hybrid promoters such as ptac (pDR540). See Brosius, et al. (1988) "Expression Vectors Employing Lambda-, trp-, lac-, and Ipp-derived Promoters", in Rodriguez and Denhardt (eds.) Vectors: A Survey of Molecular Cloning Vectors and Their Uses 10:205-236  
10 Buttersworth, Boston, MA.

Lower eukaryotes, e.g., yeasts and Dictyostelium, may be transformed with F06B09 protein sequence containing vectors. For purposes of this invention, the most common lower eukaryotic host is the baker's yeast, Saccharomyces cerevisiae. It will be used generically to represent lower eukaryotes although a number of other strains and species will be essentially equivalent. Yeast vectors typically consist of a replication origin (unless of the integrating type), one or more selection genes, a promoter, DNA encoding the desired protein or its fragments, and sequences for translation termination, polyadenylation, and transcription termination. Suitable expression vectors for yeast include such constitutive promoters as 3-phosphoglycerate kinase and various other glycolytic enzyme gene promoters or such inducible promoters as the alcohol dehydrogenase 2 promoter or metallothionine promoter. Suitable vectors include derivatives of the following types: self-replicating low copy number (such as the YRp-series), self-replicating high copy number (such as the YEpl-series), integrating types (such as the YIp-series), or mini-chromosomes (such as the YCp-series).  
15  
20  
25  
30

Higher eukaryotic tissue culture cells are typically the preferred host cells for expression of the functionally active F06B09 protease polypeptides. In principle, many higher eukaryotic tissue culture cell lines may be used, e.g., insect baculovirus expression systems, whether from an invertebrate or vertebrate source. However, mammalian cells are preferred to  
35

achieve proper natural processing, both cotranslationally and posttranslationally. Transformation or transfection and propagation of such cells are routine. Useful cell lines include HeLa cells, Chinese hamster ovary (CHO) cell lines, baby rat kidney (BRK) cell lines, insect cell lines, bird cell lines, and monkey (COS) cell lines. Expression vectors for such cell lines usually include an origin of replication, a promoter, a translation initiation site, RNA splice sites (e.g., if genomic DNA is used), a polyadenylation site, and a transcription termination site. These vectors also may contain selection and/or amplification genes. Suitable expression vectors may be plasmids, viruses, or retroviruses carrying promoters derived, e.g., from such sources as from adenovirus, SV40, parvoviruses, vaccinia virus, or cytomegalovirus. Representative examples of suitable expression vectors include pCDNA1; pCD, see Okayama, et al. (1985) Mol. Cell Biol. 5:1136-1142; pMC1neo Poly-A, see Thomas, et al. (1987) Cell 51:503-512; and a baculovirus vector such as pAC 373 or pAC 610.

It is likely that F06B09 protein need not be glycosylated to elicit biological responses. However, it will occasionally be desirable to express an F06B09 polypeptide in a system which provides a specific or defined glycosylation pattern. In this case, the usual pattern will be that provided naturally by the expression system. However, the pattern will be modifiable by exposing the polypeptide, e.g., in unglycosylated form, to appropriate glycosylating proteins introduced into a heterologous expression system. For example, an F06B09 protein gene may be co-transformed with one or more genes encoding mammalian or other glycosylating enzymes. It is further understood that over glycosylation may be detrimental to F06B09 protein biological activity, and that one of skill may perform routine testing to optimize the degree of glycosylation which confers optimal biological activity.

An F06B09 protein, or a fragment thereof, may be engineered to be phosphatidyl inositol (PI) linked to a

cell membrane, but can be removed from membranes by treatment with a phosphatidyl inositol cleaving enzyme, e.g., phosphatidyl inositol phospholipase-C. This releases the antigen in a biologically active form, and allows purification by standard procedures of protein chemistry. See, e.g., Low (1989) Biochem. Biophys. Acta 988:427-454; Tse, et al. (1985) Science 230:1003-1008; and Brunner, et al. (1991) J. Cell Biol. 114:1275-1283.

Now that F06B09 proteins have been characterized, fragments or derivatives thereof can be prepared by conventional processes for synthesizing peptides. These include processes such as are described in Stewart and Young (1984) Solid Phase Peptide Synthesis Pierce Chemical Co., Rockford, IL; Bodanszky and Bodanszky (1984) The Practice of Peptide Synthesis Springer-Verlag, New York, NY; Bodanszky (1984) The Principles of Peptide Synthesis Springer-Verlag, New York, NY; and Dawson, et al. (1994) Science 266:776-779. For example, an azide process, an acid chloride process, an acid anhydride process, a mixed anhydride process, an active ester process (for example, p-nitrophenyl ester, N-hydroxysuccinimide ester, or cyanomethyl ester), a carbodiimidazole process, an oxidative-reductive process, or a dicyclohexylcarbodiimide (DCCD)/additive process can be used. Solid phase and solution phase syntheses are both applicable to the foregoing processes.

The prepared protein and fragments thereof can be isolated and purified from the reaction mixture by means of peptide separation, for example, by extraction, precipitation, electrophoresis and various forms of chromatography, and the like. The F06B09 proteins of this invention can be obtained in varying degrees of purity depending upon its desired use. Purification can be accomplished by use of known protein purification techniques or by the use of the antibodies or binding partners herein described, e.g., in immunoabsorbant affinity chromatography. This immunoabsorbant affinity chromatography is carried out, e.g., by first linking the antibodies to a solid support and then contacting the

linked antibodies with solubilized lysates of appropriate source cells, lysates of other cells expressing the ligand, or lysates or supernatants of cells producing the F06B09 proteins as a result of recombinant DNA

5 techniques, see below.

Multiple cell lines may be screened for one which expresses an F06B09 polypeptide or protein at a high level compared with other cells. Various cell lines, e.g., a mouse thymic stromal cell line TA4, is screened  
10 and selected for its favorable handling properties. Natural F06B09 proteins can be isolated from natural sources, or by expression from a transformed cell using an appropriate expression vector. Purification of the expressed protein is achieved by standard procedures, or  
15 may be combined with engineered means for effective purification at high efficiency from cell lysates or supernatants. Epitope or other tags, e.g., FLAG or His<sub>6</sub> segments, can be used for such purification features.

## 20 V. Antibodies

Antibodies can be raised to various F06B09 proteins, including individual, polymorphic, allelic, strain, or species variants, and fragments thereof, both in their naturally occurring (full-length) forms and in their  
25 recombinant forms. Additionally, antibodies can be raised to F06B09 proteins in either their native (or active) forms or in their inactive, e.g., denatured, forms. Anti-idiotypic antibodies may also be used.

### A. Antibody Production

30 A number of immunogens may be used to produce antibodies specifically reactive with F06B09 proteins. Recombinant protein is a preferred immunogen for the production of monoclonal or polyclonal antibodies. Naturally occurring protein may also be used either in  
35 pure or impure form. Synthetic peptides, made using the human F06B09 protein sequences described herein, may also be used as an immunogen for the production of antibodies to F06B09 proteins. Recombinant protein can be expressed in eukaryotic or prokaryotic cells as described herein, and

purified as described. Naturally folded or denatured material can be used, as appropriate, for producing antibodies. Either monoclonal or polyclonal antibodies may be generated, e.g., for subsequent use in  
5 immunoassays to measure the protein.

Methods of producing polyclonal antibodies are well known to those of skill in the art. Typically, an immunogen, preferably a purified protein, is mixed with an adjuvant and animals are immunized with the mixture.  
10 The animal's immune response to the immunogen preparation is monitored by taking test bleeds and determining the titer of reactivity to the F06B09 protein of interest. For example, when appropriately high titers of antibody to the immunogen are obtained, usually after repeated  
15 immunizations, blood is collected from the animal and antisera are prepared. Further fractionation of the antisera to enrich for antibodies reactive to the protein can be done if desired. See, e.g., Harlow and Lane; or Coligan.

20 Monoclonal antibodies may be obtained by various techniques familiar to those skilled in the art. Typically, spleen cells from an animal immunized with a desired antigen are immortalized, commonly by fusion with a myeloma cell (see, Kohler and Milstein (1976) Eur. J. Immunol. 6:511-519, incorporated herein by reference).  
25 Alternative methods of immortalization include transformation with Epstein Barr Virus, oncogenes, or retroviruses, or other methods known in the art. Colonies arising from single immortalized cells are  
30 screened for production of antibodies of the desired specificity and affinity for the antigen, and yield of the monoclonal antibodies produced by such cells may be enhanced by various techniques, including injection into the peritoneal cavity of a vertebrate host.  
35 Alternatively, one may isolate DNA sequences which encode a monoclonal antibody or a binding fragment thereof by screening a DNA library from human B cells according, e.g., to the general protocol outlined by Huse, et al. (1989) Science 246:1275-1281.

Antibodies, including binding fragments and single chain versions, against predetermined fragments of F06B09 proteins can be raised by immunization of animals with conjugates of the fragments with carrier proteins as described above. Monoclonal antibodies are prepared from cells secreting the desired antibody. These antibodies can be screened for binding to normal or defective F06B09 protein, or screened for agonistic or antagonistic activity, e.g., mediated through a receptor. These monoclonal antibodies will usually bind with at least a  $K_D$  of about 1 mM, more usually at least about 300  $\mu$ M, typically at least about 100  $\mu$ M, more typically at least about 30  $\mu$ M, preferably at least about 10  $\mu$ M, and more preferably at least about 3  $\mu$ M or better.

In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts, such as mice, rodents, primates, humans, etc. Description of techniques for preparing such monoclonal antibodies may be found in, e.g., Stites, et al. (eds.) Basic and Clinical Immunology (4th ed.) Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane (1988) Antibodies: A Laboratory Manual CSH Press; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed.) Academic Press, New York, NY; and particularly in Kohler and Milstein (1975) Nature 256:495-497, which discusses one method of generating monoclonal antibodies. Summarized briefly, this method involves injecting an animal with an immunogen. The animal is then sacrificed and cells taken from its spleen, which are then fused with myeloma cells. The result is a hybrid cell or "hybridoma" that is capable of reproducing in vitro. The population of hybridomas is then screened to isolate individual clones, each of which secrete a single antibody species to the immunogen. In this manner, the individual antibody species obtained are the products of immortalized and cloned single B cells from the immune animal generated in response to a specific site recognized on the immunogenic substance.

Other suitable techniques involve selection of libraries of antibodies in phage or similar vectors. See, e.g., Huse, et al. (1989) "Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage Lambda," Science 246:1275-1281; and Ward, et al. (1989) Nature 341:544-546. The polypeptides and antibodies of the present invention may be used with or without modification, including chimeric or humanized antibodies. Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic particles, and the like. Patents teaching the use of such labels include U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant immunoglobulins may be produced, see, Cabilly, U.S. Patent No. 4,816,567; and Queen, et al. (1989) Proc. Nat'l Acad. Sci. USA 86:10029-10033; or made in transgenic mice, see Mendez, et al. (1997) Nature Genetics 15:146-156.

The antibodies of this invention are useful for affinity chromatography in isolating F06B09 protein. Columns can be prepared where the antibodies are linked to a solid support, e.g., particles, such as agarose, SEPHADEX, or the like, where a cell lysate or supernatant may be passed through the column, the column washed, followed by increasing concentrations of a mild denaturant, whereby purified F06B09 protein will be released. The converse can be performed using protein to isolate specific antibodies.

Other antibodies may block enzymatic activity. The antibodies may also be used to screen expression libraries for particular expression products. Usually the antibodies used in such a procedure will be labeled

with a moiety allowing easy detection of presence of antigen by antibody binding.

Antibodies to F06B09 proteins may be used for the identification of cell populations expressing F06B09 protein. By assaying the expression products of cells expressing F06B09 proteins it is possible to diagnose disease, e.g., metabolic conditions. The proteins may also be markers for specific tissue or cell subpopulations, e.g., dendritic cells.

Antibodies raised against each F06B09 protein will also be useful to raise anti-idiotypic antibodies. These will be useful in detecting or diagnosing various immunological conditions related to expression of the respective antigens.

15

#### B. Immunoassays

A particular protein can be measured by a variety of immunoassay methods. For a review of immunological and immunoassay procedures in general, see Stites and Terr (eds. 1991) Basic and Clinical Immunology (7th ed.). Moreover, the immunoassays of the present invention can be performed in many configurations, which are reviewed extensively in Maggio (ed. 1980) Enzyme Immunoassay CRC Press, Boca Raton, Florida; Tijan (1985) "Practice and Theory of Enzyme Immunoassays," Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Publishers B.V., Amsterdam; and Harlow and Lane Antibodies, A Laboratory Manual, supra, each of which is incorporated herein by reference. See also Chan (ed. 1987) Immunoassay: A Practical Guide Academic Press, Orlando, FL; Price and Newman (eds. 1991) Principles and Practice of Immunoassays Stockton Press, NY; and Ngo (ed. 1988) Non-isotopic Immunoassays Plenum Press, NY.

Immunoassays for measurement of F06B09 proteins or peptides can be performed by a variety of methods known to those skilled in the art. In brief, immunoassays to measure the protein can be either competitive or noncompetitive binding assays. In competitive binding assays, the sample to be analyzed competes with a labeled



analyte for specific binding sites on a capture agent bound to a solid surface. Preferably the capture agent is an antibody specifically reactive with F06B09 proteins produced as described above. The concentration of  
5 labeled analyte bound to the capture agent is inversely proportional to the amount of free analyte present in the sample.

In a competitive binding immunoassay, the F06B09 protein present in the sample competes with labeled  
10 protein for binding to a specific binding agent, for example, an antibody specifically reactive with the F06B09 protein. The binding agent may be bound to a solid surface to effect separation of bound labeled protein from the unbound labeled protein. Alternately,  
15 the competitive binding assay may be conducted in liquid phase and a variety of techniques known in the art may be used to separate the bound labeled protein from the unbound labeled protein. Following separation, the amount of bound labeled protein is determined. The  
20 amount of protein present in the sample is inversely proportional to the amount of labeled protein binding.

Alternatively, a homogeneous immunoassay may be performed in which a separation step is not needed. In these immunoassays, the label on the protein is altered  
25 by the binding of the protein to its specific binding agent. This alteration in the labeled protein results in a decrease or increase in the signal emitted by label, so that measurement of the label at the end of the immunoassay allows for detection or quantitation of the  
30 protein.

F06B09 proteins may also be determined by a variety of noncompetitive immunoassay methods. For example, a two-site, solid phase sandwich immunoassay may be used. In this type of assay, a binding agent for the protein,  
35 for example an antibody, is attached to a solid support. A second protein binding agent, which may also be an antibody, and which binds the protein at a different site, is labeled. After binding at both sites on the protein has occurred, the unbound labeled binding agent

is removed and the amount of labeled binding agent bound to the solid phase is measured. The amount of labeled binding agent bound is directly proportional to the amount of protein in the sample.

5 Western blot analysis can be used to determine the presence of F06B09 proteins in a sample. Electrophoresis is carried out, for example, on a tissue sample suspected of containing the protein. Following electrophoresis to separate the proteins, and transfer of the proteins to a  
10 suitable solid support, e.g., a nitrocellulose filter, the solid support is incubated with an antibody reactive with the protein. This antibody may be labeled, or alternatively may be detected by subsequent incubation with a second labeled antibody that binds the primary  
15 antibody.

The immunoassay formats described above may employ labeled assay components. The label may be coupled directly or indirectly to the desired component of the assay according to methods well known in the art. A wide  
20 variety of labels and methods may be used. Traditionally, a radioactive label incorporating  $^3\text{H}$ ,  $^{125}\text{I}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ , or  $^{32}\text{P}$  was used. Non-radioactive labels include ligands which bind to labeled antibodies, fluorophores, chemiluminescent agents, enzymes, and  
25 antibodies which can serve as specific binding pair members for a labeled ligand. The choice of label depends on sensitivity required, ease of conjugation with the compound, stability requirements, and available instrumentation. For a review of various labeling or  
30 signal producing systems which may be used, see U.S. Patent No. 4,391,904, which is incorporated herein by reference.

Antibodies reactive with a particular protein can also be measured by a variety of immunoassay methods.  
35 For a review of immunological and immunoassay procedures applicable to the measurement of antibodies by immunoassay techniques, see Stites and Terr (eds.) Basic and Clinical Immunology (7th ed.) supra; Maggio (ed.)

Enzyme Immunoassay, supra; and Harlow and Lane  
Antibodies, A Laboratory Manual, supra.

In brief, immunoassays to measure antisera reactive with F06B09 proteins can be either competitive or noncompetitive binding assays. In competitive binding assays, the sample analyte competes with a labeled analyte for specific binding sites on a capture agent bound to a solid surface. Preferably the capture agent is a purified recombinant F06B09 protein produced as described above. Other sources of F06B09 proteins, including isolated or partially purified naturally occurring protein, may also be used. Noncompetitive assays include sandwich assays, in which the sample analyte is bound between two analyte-specific binding reagents. One of the binding agents is used as a capture agent and is bound to a solid surface. The second binding agent is labeled and is used to measure or detect the resultant complex by visual or instrument means. A number of combinations of capture agent and labeled binding agent can be used. A variety of different immunoassay formats, separation techniques, and labels can be also be used similar to those described above for the measurement of F06B09 proteins. Similar methods may be used to evaluate or quantitate specific binding compounds.

#### VI. Purified F06B09 proteins

Human F06B09 protein amino acid sequences are provided in Table 1 and SEQ ID NO: 2 or 4.

Purified protein or defined peptides are useful for generating antibodies by standard methods, as described above. Synthetic peptides or purified protein can be presented to an immune system to generate polyclonal and monoclonal antibodies. See, e.g., Coligan (1991) Current Protocols in Immunology Wiley/Greene, NY; and Harlow and Lane (1989) Antibodies: A Laboratory Manual Cold Spring Harbor Press, NY, which are incorporated herein by reference.

The specific binding composition can be used for screening an expression library made from a cell line which expresses an F06B09 protein. Many methods for screening are available, e.g., standard staining of surface expressed ligand, or by panning. Screening of intracellular expression can also be performed by various staining or immunofluorescence procedures. The binding compositions could be used to affinity purify or sort out cells expressing the ligand.

The peptide segments, along with comparison to homologous genes, can also be used to produce appropriate oligonucleotides to screen a library. The genetic code can be used to select appropriate oligonucleotides useful as probes for screening. In combination with polymerase chain reaction (PCR) techniques, synthetic oligonucleotides will be useful in selecting desired clones from a library, including natural allelic and polymorphic variants.

The peptide sequences allow preparation of peptides to generate antibodies to recognize such segments, and allow preparation of oligonucleotides which encode such sequences. The sequence also allows for synthetic preparation, e.g., see Dawson, et al. (1994) Science 266:776-779. Analysis of the structural features in comparison with the most closely related reported sequences has revealed similarities with other proteins, particularly the class of proteins known as proteases.

#### VII. Physical Variants

This invention also encompasses proteins or peptides having substantial amino acid sequence similarity with an amino acid sequence of an F06B09 protein. Natural variants include individual, polymorphic, allelic, strain, or species variants. Conservative substitutions in the amino acid sequence will normally preserve most relevant biological activities. In particular, various substitutions can be made, e.g., embodiments with 10-fold substitutions, 7-fold substitutions, 5-fold substitutions, 3-fold substitutions, 2-fold, and etc.

Such embodiments will typically retain particular features, e.g., antigenicity, with the natural forms.

Amino acid sequence similarity, or sequence identity, is determined by optimizing residue matches, if  
5 necessary, by introducing gaps as required. This changes when considering conservative substitutions as matches. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid,  
10 glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. Homologous amino acid sequences include natural polymorphic, allelic, and interspecies variations in each respective protein sequence. Typical homologous proteins  
15 or peptides will have from 50-100% similarity (if gaps can be introduced), to 75-100% similarity (if conservative substitutions are included) with the amino acid sequence of the F06B09 protein. Similarity measures will be at least about 50%, generally at least about 60%,  
20 more generally at least about 65%, usually at least about 70%, more usually at least about 75%, preferably at least about 80%, and more preferably at least about 80%, and in particularly preferred embodiments, at least about 85% or more. See also Needleham, et al. (1970) J. Mol. Biol.  
25 48:443-453; Sankoff, et al. (1983) Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison Chapter One, Addison-Wesley, Reading, MA; and software packages from IntelliGenetics, Mountain View, CA; and the University of Wisconsin Genetics  
30 Computer Group, Madison, WI.

Natural nucleic acids encoding mammalian F06B09 proteins will typically hybridize to the nucleic acid sequence of SEQ ID NO: 1 or 3 under stringent conditions. For example, nucleic acids encoding human F06B09 proteins  
35 will normally hybridize to the nucleic acid of SEQ ID NO: 1 or 3 under stringent hybridization conditions. Generally, stringent conditions are selected to be about 10° C lower than the thermal melting point (T<sub>m</sub>) for the probe sequence at a defined ionic strength and pH. The

T<sub>m</sub> is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is about  
5 0.2 molar at pH 7 and the temperature is at least about 50° C. Other factors may significantly affect the stringency of hybridization, including, among others, base composition and size of the complementary strands, the presence of organic solvents such as formamide, and  
10 the extent of base mismatching. A preferred embodiment will include nucleic acids which will bind to disclosed sequences in 50% formamide and 200 mM NaCl at 42° C. See, e.g., Sambrook, et al.

An isolated F06B09 protein DNA can be readily  
15 modified by nucleotide substitutions, nucleotide deletions, nucleotide insertions, and short inversions of nucleotide stretches. These modifications result in novel DNA sequences which encode F06B09 protein antigens, their derivatives, or proteins having highly similar  
20 physiological, immunogenic, or antigenic activity.

Modified sequences can be used to produce mutant antigens or to enhance expression. Enhanced expression may involve gene amplification, increased transcription, increased translation, and other mechanisms. Such mutant  
25 F06B09 protein derivatives include predetermined or site-specific mutations of the respective protein or its fragments. "Mutant F06B09 protein" encompasses a polypeptide otherwise falling within the homology definition of the human F06B09 protein as set forth  
30 above, but having an amino acid sequence which differs from that of an F06B09 protein as found in nature, whether by way of deletion, substitution, or insertion. In particular, "site specific mutant F06B09 protein" generally includes proteins having significant similarity  
35 with a protein having a sequence of SEQ ID NO: 2 or 4, and as sharing various biological activities, e.g., antigenic or immunogenic, with those sequences, and in preferred embodiments contain most or all of the disclosed sequence. This applies also to polymorphic

variants from different individuals. Similar concepts apply to different F06B09 proteins, particularly those found in various warm blooded animals, e.g., mammals and birds. As stated before, it is emphasized that

5 descriptions are generally meant to encompass other F06B09 proteins, not limited to the human embodiments specifically discussed.

Although site specific mutation sites are predetermined, mutants need not be site specific. F06B09

10 protein mutagenesis can be conducted by making amino acid insertions or deletions. Substitutions, deletions, insertions, or combinations may be generated to arrive at a final construct. Insertions include amino- or carboxyl- terminal fusions, e.g. epitope tags. Random

15 mutagenesis can be conducted at a target codon and the expressed mutants can then be screened for the desired activity. Methods for making substitution mutations at predetermined sites in DNA having a known sequence are well known in the art, e.g., by M13 primer mutagenesis or

20 polymerase chain reaction (PCR) techniques. See also, Sambrook, et al. (1989) and Ausubel, et al. (1987 and Supplements). The mutations in the DNA normally should not place coding sequences out of reading frames and preferably will not create complementary regions that

25 could hybridize to produce secondary mRNA structure such as loops or hairpins.

The present invention also provides recombinant proteins, e.g., heterologous fusion proteins using segments from these proteins. A heterologous fusion

30 protein is a fusion of proteins or segments which are naturally not normally fused in the same manner. Thus, the fusion product of an immunoglobulin with an F06B09 protein polypeptide is a continuous protein molecule having sequences fused in a typical peptide linkage,

35 typically made as a single translation product and exhibiting properties derived from each source peptide. A similar concept applies to heterologous nucleic acid sequences. Antibody fusion proteins are also contemplated.

In addition, new constructs may be made from combining similar functional domains from other proteins. For example, protein-binding or other segments may be "swapped" between different new fusion polypeptides or fragments. See, e.g., Cunningham, et al. (1989) Science 243:1330-1336; and O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992. Thus, new chimeric polypeptides exhibiting new combinations of specificities will result from the functional linkage of protein-binding specificities and other functional domains.

#### VIII. Binding Agent:F06B09 Protein Complexes

An F06B09 protein that specifically binds to or that is specifically immunoreactive with an antibody generated against a defined immunogen, such as an immunogen consisting of the amino acid sequence of SEQ ID NO: 2 or 4, is typically determined in an immunoassay. The immunoassay uses a polyclonal antiserum which was raised to a protein of SEQ ID NO: 2 or 4. This antiserum is selected to have low crossreactivity against other proteases and any such crossreactivity is removed by immunoabsorption prior to use in the immunoassay.

In order to produce antisera for use in an immunoassay, the protein of SEQ ID NO: 2 or 4 is isolated as described herein. For example, recombinant protein may be produced in a mammalian cell line. An inbred strain of mice such as Balb/c is immunized with the protein of SEQ ID NO: 2 or 4 using a standard adjuvant, such as Freund's adjuvant, and a standard mouse immunization protocol (see Harlow and Lane, supra). Alternatively, a synthetic peptide, preferably near full length, derived from the sequences disclosed herein and conjugated to a carrier protein can be used an immunogen. Polyclonal sera are collected and titered against the immunogen protein in an immunoassay, for example, a solid phase immunoassay with the immunogen immobilized on a solid support. Polyclonal antisera with a titer of  $10^4$  or greater are selected and tested for their cross reactivity against other proteases, e.g., using a



competitive binding immunoassay such as the one described in Harlow and Lane, supra, at pages 570-573. Preferably, two related proteins are used in this determination in conjunction with either F06B09 protein.

5           Immunoassays in the competitive binding format can be used for the crossreactivity determinations. For example, a protein of SEQ ID NO: 4 can be immobilized to a solid support. Proteins added to the assay compete with the binding of the antisera to the immobilized  
10 antigen. The ability of the above proteins to compete with the binding of the antisera to the immobilized protein is compared to the protein of SEQ ID NO: 4. The percent crossreactivity for the above proteins is calculated, using standard calculations. Those antisera  
15 with less than 10% crossreactivity with each of the proteins listed above are selected and pooled. The cross-reacting antibodies are then removed from the pooled antisera by immunoabsorption with the above-listed proteins.

20           The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay as described above to compare a second protein to the immunogen protein (e.g., the protein motif of SEQ ID NO: 4). In order to make this comparison, the two proteins are each assayed  
25 at a wide range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein required is less than twice the amount of the protein of SEQ ID NO: 4 that is  
30 required, then the second protein is said to specifically bind to an antibody generated to the immunogen.

          It is understood that F06B09 proteins are families of homologous proteins that comprise two or more genes. For a particular gene product, such as the human F06B09  
35 proteins, the term refers not only to the amino acid sequences disclosed herein, but also to other proteins that are polymorphic, allelic, non-allelic, or species variants or equivalents. It is also understood that the term "human F06B09 protein" includes equivalent proteins,

e.g., nonnatural mutations introduced by deliberate mutation using conventional recombinant technology such as single site mutation, or by excising short sections of DNA encoding F06B09 proteins, or by substituting new amino acids, or adding new amino acids. Such minor alterations must substantially maintain the immunoidentity of the original molecule and/or its biological activity. Thus, these alterations include proteins that are specifically immunoreactive with a designated naturally occurring F06B09 protein, for example, the human F06B09 protein shown in SEQ ID NO: 4. The biological properties of the altered proteins can be determined by expressing the protein in an appropriate cell line and measuring, e.g., enzymatic activity under appropriate conditions. Particular protein modifications considered minor would include conservative substitution of amino acids with similar chemical properties, as described above for F06B09 protein families as a whole. By aligning a protein optimally with the protein of SEQ ID NO: 4, and by using the conventional immunoassays described herein to determine immunoidentity, or by using lymphocyte chemotaxis assays, one can determine the protein compositions of the invention.

## IX. Functional Variants

The blocking of physiological response to F06B09 protein may result from the inhibition of enzymatic activity of the protein against its substrate, e.g., through competitive inhibition. Thus, in vitro assays of the present invention will often use isolated protein, membranes from cells expressing a recombinant membrane associated proteins, soluble fragments comprising enzymatically active segments of these proteins, or fragments attached to solid phase substrates. These assays will also allow for the diagnostic determination of the effects of either binding segment mutations and modifications, or protein mutations and modifications, e.g., protein analogs. This invention also contemplates the use of competitive drug screening assays, e.g., where

neutralizing antibodies to antigen or enzyme fragments compete with a test compound for binding to the protein. In this manner, the antibodies can be used to detect the presence of a polypeptide which shares one or more  
5 antigenic binding sites of the protein and can also be used to occupy binding sites on the protein that might otherwise interact with, e.g., substrate.

"Derivatives" of F06B09 proteins include amino acid sequence mutants, glycosylation variants, and covalent or  
10 aggregate conjugates with other chemical moieties. Covalent derivatives can be prepared by linkage of functionalities to groups which are found in F06B09 protein amino acid side chains or at the N- or C-termini, e.g., by means which are well known in the art.  
15 These derivatives can include, without limitation, aliphatic esters or amides of the carboxyl terminus, or of residues containing carboxyl side chains, O-acyl derivatives of hydroxyl group-containing residues, and N-acyl derivatives of the amino terminal amino acid or  
20 amino-group containing residues, e.g., lysine or arginine. Acyl groups are typically selected from the group of alkyl-moieties including, e.g., C3 to C18 normal alkyl, thereby forming alkanoyl aroyl species. Covalent attachment to carrier proteins may be important when  
25 immunogenic moieties are haptens.

In particular, glycosylation alterations are included, e.g., made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing, or in further processing steps. Particularly  
30 preferred means for accomplishing this are by exposing the polypeptide to glycosylating enzymes derived from cells which normally provide such processing, e.g., mammalian glycosylation enzymes. Deglycosylation enzymes are also contemplated. Also embraced are versions of the  
35 same primary amino acid sequence which have other minor modifications, including phosphorylated amino acid residues, e.g., phosphotyrosine, phosphoserine, or phosphothreonine, or other moieties, including ribosyl groups or cross-linking reagents.

A major group of derivatives are covalent conjugates of the F06B09 protein or fragments thereof with other proteins or polypeptides. These derivatives can be synthesized in recombinant culture such as N- or C-terminal fusions or by the use of agents known in the art for their usefulness in cross-linking proteins through reactive side groups. Preferred protein derivatization sites with cross-linking agents are at free amino groups, carbohydrate moieties, and cysteine residues.

10 Fusion polypeptides between human F06B09 proteins and other homologous or heterologous proteins are also provided. Heterologous polypeptides may be fusions between different related proteins, resulting in, e.g., a hybrid protein exhibiting modified substrate or other  
15 binding specificity. Likewise, heterologous fusions may be constructed which would exhibit a combination of properties or activities of the derivative proteins. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with a segment or domain of a protein, e.g., a receptor-binding segment, so that the presence or  
20 location of the fused protein may be easily determined. See, e.g., Dull, et al., U.S. Patent No. 4,859,609. Other gene fusion partners include bacterial  $\beta$ -galactosidase, trpE, Protein A,  $\beta$ -lactamase, alpha  
25 amylase, alcohol dehydrogenase, and yeast alpha mating factor. See, e.g., Godowski, et al. (1988) Science 241:812-816.

Such polypeptides may also have amino acid residues which have been chemically modified by phosphorylation, sulfonation, biotinylation, or the addition or removal of  
30 other moieties, particularly those which have molecular shapes similar to phosphate groups. In some embodiments, the modifications will be useful labeling reagents, or serve as purification targets, e.g., affinity ligands.

35 This invention also contemplates the use of derivatives of F06B09 proteins other than variations in amino acid sequence or glycosylation. Such derivatives may involve covalent or aggregative association with chemical moieties. These derivatives generally include

the three classes: (1) salts, (2) side chain and terminal residue covalent modifications, and (3) adsorption complexes, for example with cell membranes. Such covalent or aggregative derivatives are useful as immunogens, as reagents in immunoassays, or in purification methods such as for affinity purification of ligands or other binding ligands. For example, an F06B09 protein can be immobilized by covalent bonding to a solid support such as cyanogen bromide-activated SEPHAROSE, by methods which are well known in the art, or adsorbed onto polyolefin surfaces, with or without glutaraldehyde cross-linking, for use in the assay or purification of anti-F06B09 protein antibodies. The F06B09 proteins can also be labeled with a detectable group, e.g., radioiodinated by the chloramine T procedure, covalently bound to rare earth chelates, or conjugated to another fluorescent moiety for use in diagnostic assays. Purification of F06B09 proteins may be effected by immobilized antibodies or substrate.

Isolated F06B09 protein genes will allow transformation of cells lacking expression of corresponding F06B09 proteins, e.g., either species types or cells which lack corresponding proteins and exhibit negative background activity. Expression of transformed genes will allow isolation of antigenically pure cell lines, with defined or single specie variants. This approach will allow for more sensitive detection and discrimination of the physiological effects of F06B09 protein substrate proteins. Subcellular fragments, e.g., cytoplasts or membrane fragments, can be isolated and used.

#### X. Uses

The present invention provides reagents which will find use in diagnostic applications as described elsewhere herein, e.g., in the general description for metabolic abnormalities, or below in the description of kits for diagnosis.

F06B09 protein nucleotides, e.g., human F06B09 protein DNA or RNA, may be used as a component in a forensic assay. For instance, the nucleotide sequences provided may be labeled using, e.g.,  $^{32}\text{P}$  or biotin and  
5 used to probe standard restriction fragment polymorphism blots, providing a measurable character to aid in distinguishing between individuals. Such probes may be used in well-known forensic techniques such as genetic fingerprinting. In addition, nucleotide probes made from  
10 F06B09 protein sequences may be used in in situ assays to detect chromosomal abnormalities.

Antibodies and other binding agents directed towards F06B09 proteins or nucleic acids may be used to purify the corresponding F06B09 protein molecule. As described  
15 in the Examples below, antibody purification of F06B09 protein components is both possible and practicable. Antibodies and other binding agents may also be used in a diagnostic fashion to determine whether F06B09 protein components are present in a tissue sample or cell  
20 population using well-known techniques described herein. The ability to attach a binding agent to an F06B09 protein provides a means to diagnose disorders associated with F06B09 protein misregulation. Antibodies and other F06B09 protein binding agents may also be useful as  
25 histological or sorting markers. As described in the examples below, F06B09 protein expression is limited to specific tissue types. By directing a probe, such as an antibody or nucleic acid to an F06B09 protein, it is possible to use the probe to distinguish tissue and cell  
30 types in situ or in vitro.

This invention also provides reagents with significant therapeutic value. The F06B09 protein (naturally occurring or recombinant), fragments thereof, and antibodies thereto, along with compounds identified  
35 as having binding affinity to an F06B09 protein, are useful in the treatment of conditions associated with abnormal metabolism, physiology, or development, including abnormal immune responsiveness or non-responsiveness. Abnormal proliferation, regeneration,

degeneration, and atrophy may be modulated by appropriate therapeutic treatment using the compositions provided herein. The F06B09 proteins likely play roles in regulation or development of hematopoietic cells, e.g., lymphoid cells, which affect immunological responses. Thus, for example, an antagonist of an F06B09 protein could be useful in blocking the conversion of an immature or inactive immunologically relevant pro-protein to the mature or active form. Since the F06B09 proteases were derived from dendritic cells, antagonists could also be important in preventing antigen processing and/or subsequent presentation. In addition, effects on DC migration or dendrite extension between cells may result. One potential therapeutic application of F06B09 would be to block this protease in inflammatory processes involving the dendritic cells (DC). The blocking could occur on the F06B09 MMP itself or on other molecules interacting with it.

Other abnormal developmental conditions are known in cell types shown to possess F06B09 protein encoding mRNA by northern blot analysis. See Berkow (ed.) The Merck Manual of Diagnosis and Therapy, Merck & Co., Rahway, NJ; and Thorn, et al. Harrison's Principles of Internal Medicine, McGraw-Hill, NY. Developmental or functional abnormalities, e.g., of the immune system, cause significant medical abnormalities and conditions which may be susceptible to prevention or treatment using compositions provided herein.

Recombinant F06B09 protein antibodies can be purified and then administered to a patient. These reagents can be combined for therapeutic use with additional active or inert ingredients, e.g., in conventional pharmaceutically acceptable carriers or diluents, e.g., immunogenic adjuvants, along with physiologically innocuous stabilizers and excipients. These combinations can be sterile filtered and placed into dosage forms as by lyophilization in dosage vials or storage in stabilized aqueous preparations. This invention also contemplates use of antibodies or binding

fragments thereof, including forms which are not complement binding.

Drug screening using antibodies or fragments thereof can identify compounds having binding affinity to F06B09 protein, including isolation of associated components. Various substrate candidates can be screened. Subsequent biological assays can then be utilized to determine if the compound has intrinsic enzyme blocking activity. Likewise, a compound having intrinsic stimulating activity might activate the activity of an F06B09 protein. This invention further contemplates the therapeutic use of antibodies to F06B09 protein as antagonists. This approach should be particularly useful with other F06B09 protein polymorphic or species variants.

The quantities of reagents necessary for effective therapy will depend upon many different factors, including means of administration, target site, physiological state of the patient, and other medicants administered. Thus, treatment dosages should be titrated to optimize safety and efficacy. Typically, dosages used in vitro may provide useful guidance in the amounts useful for in situ administration of these reagents. Animal testing of effective doses for treatment of particular disorders will provide further predictive indication of human dosage. Various considerations are described, e.g., in Gilman, et al. (eds. 1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics (8th ed.) Pergamon Press; and (1990) Remington's Pharmaceutical Sciences (17th ed.) Mack Publishing Co., Easton, PA. Methods for administration are discussed therein and below, e.g., for oral, intravenous, intraperitoneal, or intramuscular administration, transdermal diffusion, and others. Pharmaceutically acceptable carriers will include water, saline, buffers, and other compounds described, e.g., in the Merck Index, Merck & Co., Rahway, NJ. Dosage ranges would ordinarily be expected to be in amounts lower than 1 mM concentrations, typically less than about 10  $\mu$ M



concentrations, usually less than about 100 nM, preferably less than about 10 pM (picomolar), and most preferably less than about 1 fM (femtomolar), with an appropriate carrier. Slow release formulations, or a  
5 slow release apparatus will often be utilized for continuous administration.

F06B09 proteins, fragments thereof; antibodies to it or its fragments; antagonists; and agonists, may be administered directly to the host to be treated or,  
10 depending on the size of the compounds, it may be desirable to conjugate them to carrier proteins such as ovalbumin or serum albumin prior to their administration. Therapeutic formulations may be administered in any conventional dosage formulation. While it is possible  
15 for the active ingredient to be administered alone, it is preferable to present it as a pharmaceutical formulation. Formulations typically comprise at least one active ingredient, as defined above, together with one or more acceptable carriers thereof. Each carrier should be both  
20 pharmaceutically and physiologically acceptable in the sense of being compatible with the other ingredients and not injurious to the patient. Formulations include those suitable for oral, rectal, nasal, or parenteral (including subcutaneous, intramuscular, intravenous and  
25 intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. See, e.g., Gilman, et al. (eds. 1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics  
30 (8th ed.) Pergamon Press; and (1990) Remington's Pharmaceutical Sciences (17th ed.) Mack Publishing Co., Easton, PA; Avis, et al. (eds. 1993) Pharmaceutical Dosage Forms: Parenteral Medications Dekker, NY; Lieberman, et al. (eds. 1990) Pharmaceutical Dosage  
35 Forms: Tablets Dekker, NY; and Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Disperse Systems Dekker, NY. The therapy of this invention may be combined with or used in association with other therapeutic agents.

Both the naturally occurring and the recombinant forms of the F06B09 proteins of this invention are particularly useful in kits and assay methods which are capable of screening compounds for binding activity to the proteins, including substrates or competitive inhibitors. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds in a short period. See, e.g., Fodor, et al. (1991) Science 251:767-773, and other descriptions of chemical diversity libraries, which describe means for testing of binding affinity by a plurality of compounds. The development of suitable assays can be greatly facilitated by the availability of large amounts of purified, soluble F06B09 protein as provided by this invention.

For example, antagonists or inhibitors can normally be found once the protein has been structurally defined. Testing of potential substrates or analogs is now possible upon the development of highly automated assay methods using a purified enzyme. In particular, new agonists and antagonists will be discovered by using screening techniques described herein. Of particular importance are compounds found to have a combined blockage activity for multiple F06B09 protein substrates, e.g., compounds which can serve as antagonists for polymorphic or species variants of an F06B09 protein. Inhibitors can be identified, which may be useful as therapeutic entities.

This invention is particularly useful for screening compounds by using recombinant protein in a variety of drug screening techniques. The advantages of using a recombinant protein in screening for specific inhibitors include: (a) improved renewable source of the F06B09 protein from a specific source; (b) potentially greater number of molecules per cell giving better signal to noise ratio in assays; and (c) species variant specificity (theoretically giving greater biological and disease specificity).

One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant DNA molecules expressing an F06B09 protein substrate. Cells may be isolated which express a

5 substrate in isolation from any others. Such cells, either in viable or fixed form, can be used for standard enzyme/substrate cleavage assays. See also, Parce, et al. (1989) Science 246:243-247; and Owicki, et al. (1990) Proc. Nat'l Acad. Sci. USA 87:4007-4011, which describe

10 sensitive methods to detect cellular responses. Competitive assays are particularly useful, where the cells (source of F06B09 protein) or homogenates are contacted and incubated with a labeled antibody having known binding affinity to the protein, such as  $^{125}\text{I}$ -

15 antibody, and a test sample whose binding affinity to the binding composition is being measured. The bound and free labeled binding compositions are then separated to assess the degree of antigen binding. The amount of test compound bound is inversely proportional to the amount of

20 labeled reagent binding to the known source. Any one of numerous techniques can be used to separate bound from free antigen to assess the degree of ligand binding. This separation step could typically involve a procedure such as adhesion to filters followed by washing, adhesion

25 to plastic followed by washing, or centrifugation of the cell membranes. Viable cells could also be used to screen for the effects of drugs on F06B09 protein mediated functions, e.g., proprotein activation, substrate cleavage, and others. Some detection methods

30 allow for elimination of a separation step, e.g., a proximity sensitive detection system.

Another method utilizes solubilized, unpurified or solubilized, purified F06B09 protein from transformed eukaryotic or prokaryotic host cells. This allows for a

35 "molecular" binding assay with the advantages of increased specificity, the ability to automate, and high drug test throughput.

Another technique for drug screening involves an approach which provides high throughput screening for

compounds having suitable binding affinity to an F06B09 protein, e.g., an antibody, is described in detail in Geysen, European Patent Application 84/03564, published on September 13, 1984. First, large numbers of different  
5 small peptide test compounds are synthesized on a solid substrate, e.g., plastic pins or some other appropriate surface, see Fodor, et al., supra. Then all the pins are reacted with solubilized, unpurified or solubilized, purified F06B09 protein antibody, and washed. The next  
10 step involves detecting bound F06B09 protein antibody.

Rational drug design may also be based upon structural studies of the molecular shapes of the F06B09 protein and other effectors or analogs. See, e.g., Methods in Enzymology vols. 202 and 203. Effectors may  
15 be other proteins which mediate other functions in response to antigen binding, or other proteins which normally interact with the substrate. One means for determining which sites interact with specific other proteins is a physical structure determination, e.g., x-  
20 ray crystallography or 2 dimensional NMR techniques. These will provide guidance as to which amino acid residues form molecular contact regions. For a detailed description of protein structural determination, see, e.g., Blundell and Johnson (1976) Protein Crystallography  
25 Academic Press, NY.

A purified F06B09 protein can be coated directly onto plates for use in the aforementioned drug screening techniques. However, non-neutralizing antibodies to these antigens can be used as capture antibodies to  
30 immobilize the respective antigen on the solid phase. Candidates for screening include for hybridomas, to find clones with desired binding specificity, or for inhibitors, e.g., of enzymatic activity.

#### 35 XI. Kits

This invention also contemplates use of F06B09 proteins, fragments thereof, peptides, and their fusion products in a variety of diagnostic kits and methods for detecting the presence of F06B09 protein or an F06B09

protein substrate. Typically the kit will have a compartment containing either a defined F06B09 peptide or gene segment or a reagent which recognizes one or the other, e.g., substrates or antibodies.

5           A kit for determining the binding affinity of a test compound to an F06B09 protein would typically comprise a test compound; a labeled compound, e.g., an antibody having known binding affinity for the F06B09 protein; a source of F06B09 protein (naturally occurring or  
10 recombinant); and a means for separating bound from free labeled compound, such as a solid phase for immobilizing the F06B09 protein. Once compounds are screened, those having suitable binding affinity to the F06B09 protein can be evaluated in suitable biological assays, as are  
15 well known in the art, to determine whether they act as agonists or antagonists to a substrate. The availability of recombinant F06B09 polypeptides also provide well defined standards for calibrating such assays.

          A preferred kit for determining the concentration  
20 of, for example, an F06B09 protein in a sample would typically comprise a labeled compound, e.g., antibody, having known binding affinity for the F06B09 protein, a source of F06B09 protein (naturally occurring or recombinant), and a means for separating the bound from  
25 free labeled compound, for example, a solid phase for immobilizing the F06B09 protein. Compartments containing reagents, and instructions, will normally be provided.

          Antibodies, including antigen binding fragments, specific for the F06B09 protein, or fragments thereof,  
30 are useful in diagnostic applications to detect the presence of elevated levels of F06B09 protein and/or its fragments. Such diagnostic assays can employ lysates, live cells, fixed cells, immunofluorescence, cell cultures, body fluids, and further can involve the  
35 detection of antigens related to the ligand in serum, or the like. Diagnostic assays may be homogeneous (without a separation step between free reagent and F06B09 protein complex) or heterogeneous (with a separation step). Various commercial assays exist, such as radioimmunoassay

(RIA), enzyme-linked immunosorbent assay (ELISA), enzyme immunoassay (EIA), enzyme-multiplied immunoassay technique (EMIT), substrate-labeled fluorescent immunoassay (SLFIA), and the like. For example, unlabeled antibodies can be employed by using a second antibody which is labeled and which recognizes the antibody to an F06B09 protein or to a particular fragment thereof. Similar assays have also been extensively discussed in the literature. See, e.g., Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH Press, NY; Chan (ed. 1987) Immunoassay: A Practical Guide Academic Press, Orlando, FL; Price and Newman (eds. 1991) Principles and Practice of Immunoassay Stockton Press, NY; and Ngo (ed. 1988) Nonisotopic Immunoassay Plenum Press, NY.

Anti-idiotypic antibodies may have similar use to diagnose presence of antibodies against an F06B09 protein, as such may be diagnostic of various abnormal states. For example, overproduction of F06B09 protein may result in production of various immunological or other medical reactions which may be diagnostic of abnormal physiological states, e.g., in cell growth, activation, or differentiation.

Frequently, the reagents for diagnostic assays are supplied in kits, so as to optimize the sensitivity of the assay. For the subject invention, depending upon the nature of the assay, the protocol, and the label, either labeled or unlabeled antibody, or labeled F06B09 protein is provided. This is usually in conjunction with other additives, such as buffers, stabilizers, materials necessary for signal production such as substrates for enzymes, and the like. Preferably, the kit will also contain instructions for proper use and disposal of the contents after use. Typically the kit has compartments for each useful reagent. Desirably, the reagents are provided as a dry lyophilized powder, where the reagents may be reconstituted in an aqueous medium providing appropriate concentrations of reagents for performing the assay.

Many of the aforementioned constituents of the drug screening and the diagnostic assays may be used without modification, or may be modified in a variety of ways. For example, labeling may be achieved by covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In any of these assays, the protein, test compound, F06B09 protein, or antibodies thereto can be labeled either directly or indirectly. Possibilities for direct labeling include label groups: radiolabels such as  $^{125}\text{I}$ , enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups.

There are also numerous methods of separating the bound from the free antigen, or alternatively the bound from the free test compound. The F06B09 protein can be immobilized on various matrices followed by washing. Suitable matrices include plastic such as an ELISA plate, filters, and beads. Methods of immobilizing the F06B09 protein to a matrix include, without limitation, direct adhesion to plastic, use of a capture antibody, chemical coupling, and biotin-avidin. The last step in this approach usually involves the precipitation of enzyme/antibody or enzyme substrate complex by various methods including those utilizing, e.g., an organic solvent such as polyethylene glycol or a salt such as ammonium sulfate. Other suitable separation techniques include, without limitation, the fluorescein antibody magnetizable particle method described in Rattle, et al. (1984) Clin. Chem. 30:1457-1461, and the double antibody magnetic particle separation as described in U.S. Pat. No. 4,659,678.

Methods for linking proteins or their fragments to the various labels have been extensively reported in the

literature and do not require detailed discussion here. Many of the techniques involve the use of activated carboxyl groups either through the use of carbodiimide or active esters to form peptide bonds, the formation of thioethers by reaction of a mercapto group with an activated halogen such as chloroacetyl, or an activated olefin such as maleimide, for linkage, or the like. Fusion proteins will also find use in these applications.

Another diagnostic aspect of this invention involves use of oligonucleotide or polynucleotide sequences taken from the sequence of an F06B09 protein. These sequences can be used as probes for detecting levels of the F06B09 protein message in samples from natural sources, or patients suspected of having an abnormal condition, e.g., immune problem. The preparation of both RNA and DNA nucleotide sequences, the labeling of the sequences, and the preferred size of the sequences has received ample description and discussion in the literature. Normally an oligonucleotide probe should have at least about 14 nucleotides, usually at least about 18 nucleotides, and the polynucleotide probes may be up to several kilobases. Various detectable labels may be employed, most commonly radionuclides, particularly  $^{32}\text{P}$ . However, other techniques may also be employed, such as using biotin modified nucleotides for introduction into a polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides, fluorophores, enzymes, or the like. Alternatively, antibodies may be employed which can recognize specific duplexes, including DNA duplexes, RNA duplexes, DNA-RNA hybrid duplexes, or DNA-protein duplexes. The antibodies in turn may be labeled and the assay carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected. The use of probes to the novel anti-sense RNA may be carried out using many conventional techniques such as nucleic acid hybridization, plus and minus screening, recombinational



probing, hybrid released translation (HRT), and hybrid arrested translation (HART). This also includes amplification techniques such as polymerase chain reaction (PCR).

5       Diagnostic kits which also test for the qualitative or quantitative presence of other markers are also contemplated. Diagnosis or prognosis may depend on the combination of multiple indications used as markers. Thus, kits may test for combinations of markers. See,  
10 e.g., Viallet, et al. (1989) Progress in Growth Factor Res. 1:89-97.

### XIII. Substrate Identification

Having isolated a protease, methods exist for  
15 identifying a target substrate. For example, a candidate substrate can be contacted with an F06B09 protein in an enzymatic reaction. The resulting cleavage or product can be analyzed, e.g., using SDS-PAGE, HPLC, spectroscopy or other forms of analysis. For example, the molecular  
20 weight of a protease cleavage product should be compared against the molecular weights of the uncleaved substrate and the F06B09 protein. The successful candidate substrate will exhibit a shift to a lower molecular weight. Analysis of the substrate should determine what  
25 site specificity may exist for the enzyme under the tested conditions. Alternatively, if the protease acts by transforming an inactive substrate to the active form, the resulting activity can be assayed, e.g., by the result of the activated factor, e.g., proliferation,  
30 apoptosis, or activation of a target cell.

Sequence specificity of products may allow search through sequence databases to identify candidate proteins as physiologically natural substrates. Alternatively, the protease may be involve in antigen processing and  
35 presentation to appropriate immune cells.

The broad scope of this invention is best understood with reference to the following examples, which are not intended to limit the invention to specific embodiments.

## EXAMPLES

## I. General Methods

Many of the standard methods below are described or  
5 referenced, e.g., in Maniatis, et al. (1982) Molecular Cloning, A Laboratory Manual Cold Spring Harbor Laboratory, Cold Spring Harbor Press, NY; Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (2d ed.) Vols. 1-3, CSH Press, NY; Ausubel, et al., Biology  
10 Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and Supplements) Current Protocols in Molecular Biology Wiley/Greene, NY; Innis, et al. (eds. 1990) PCR Protocols: A Guide to Methods and Applications Academic Press, NY. Methods for protein purification  
15 include such methods as ammonium sulfate precipitation, column chromatography, electrophoresis, centrifugation, crystallization, and others. See, e.g., Ausubel, et al. (1987 and periodic supplements); Deutscher (1990) "Guide to Protein Purification," Methods in Enzymology vol. 182, and other volumes in this series; and manufacturer's  
20 literature on use of protein purification products, e.g., Pharmacia, Piscataway, NJ, or Bio-Rad, Richmond, CA. Combination with recombinant techniques allow fusion to appropriate segments (epitope tags), e.g., to a FLAG  
25 sequence or an equivalent which can be fused, e.g., via a protein-removable sequence. See, e.g., Hochuli (1989) Chemische Industrie 12:69-70; Hochuli (1990) "Purification of Recombinant Proteins with Metal Chelate Absorbent" in Setlow (ed.) Genetic Engineering, Principle and Methods 12:87-98, Plenum Press, NY; and Crowe, et al.  
30 (1992) QIAexpress: The High Level Expression & Protein Purification System QUIAGEN, Inc., Chatsworth, CA.

Standard immunological techniques are described, e.g., in Hertenberg, et al. (eds. 1996) Weir's Handbook of Experimental Immunology vols. 1-4, Blackwell Science;  
35 Coligan (1991) Current Protocols in Immunology Wiley/Greene, NY; and Methods in Enzymology volumes. 70, 73, 74, 84, 92, 93, 108, 116, 121, 132, 150, 162, and 163. Assays for neural cell biological activities are

described, e.g., in Wouterlood (ed. 1995) Neuroscience Protocols modules 10, Elsevier; Methods in Neurosciences Academic Press; and Neuromethods Humana Press, Totowa, NJ. Methodology of developmental systems is described, e.g., in Meisami (ed.) Handbook of Human Growth and Developmental Biology CRC Press; and Chrispeels (ed.) Molecular Techniques and Approaches in Developmental Biology Interscience.

FACS analyses are described in Melamed, et al. (1990) Flow Cytometry and Sorting Wiley-Liss, Inc., New York, NY; Shapiro (1988) Practical Flow Cytometry Liss, New York, NY; and Robinson, et al. (1993) Handbook of Flow Cytometry Methods Wiley-Liss, New York, NY.

15 II. Hematopoietic factors, cells, and cell lines  
rhGM-CSF (specific activity :  $2 \times 10^6$  U/mg; Schering-Plough Research Institute, Kenilworth, NJ) was used at a saturating concentration of 100 ng/ml (200 U/ml). rhTNF $\alpha$  (specific activity :  $2 \times 10^7$  U/mg; Genzyme, Boston, MA) was used at an optimal concentration of 2.5 ng/ml (50 U/ml). rhSCF (specific activity :  $4 \times 10^5$  U/mg; R&D, Abington, U.K.) and rhM-CSF (specific activity :  $2 \times 10^6$  U/mg; R&D) were used at optimal concentration of 25 ng/ml. rhG-CSF (ED<sub>50</sub>: 0.01-0.03 ng/ml R&D) was used at an optimal concentration of 25 ng/ml.

Peripheral blood mononuclear cells (PBMC) were obtained from healthy donors after Ficoll-Hypaque gradient centrifugation ( $d=1.077$ ; Eurobio, Paris, France). T cells were purified from PBMC by immunomagnetic depletion (Dyna1, Oslo, Norway) using a cocktail of mAbs (CD14, CD16, CD35, HLA-DR, (Immunotech, Marseille, France), CD19 (ascites), NKH1 (Coulter, Hialeah, FL), CD40 (mAb 89 produced in the laboratory)). The purity of CD3<sup>+</sup> T cells was higher than 95%. T cells were activated with coated anti-CD3 and soluble anti-CD28 mAbs for 3, 12 and 24 h. B cells were obtained from human tonsils as described (Liu, et al. (1996) Immunity. 4:603-613). Briefly, T cells were first depleted by

rosetting sheep red blood cells and then the residual non-B cells were removed by immunomagnetic depletion using a cocktail of mAbs (CD2, CD3, CD4, CD14, CD16, NKH1, CD35). The purity of CD19<sup>+</sup> B cells was higher than 98%. Langerhans cells were prepared from normal skin by CD1a positive selection as described (Le Varlet, et al. (1992) J. Leukoc. Biol. 51:415-420). Granulocytes were generated in vitro from CD34<sup>+</sup> progenitors in the presence of G-CSF and SCF for 12 days. Macrophages were generated in vitro by culturing human cord blood CD34<sup>+</sup> progenitors with M-CSF and SCF for 12 days (Szabolcs, et al. (1996) Blood. 87:4520-4530). Cells were unactivated or activated by PMA-ionomycin for 1 and 6 h (PMA: 1 ng/ml, Sigma, St. Louis, MO; Ionomycin: 1 µg/ml, Calbiochem, La Jolla, CA) and pooled. The TF1 (erythrocytic), Jurkat (T cell), MRC5 (fibroblastic), JY (lymphoblastoid B cell), and U937 (myelomonocytic) cell lines were obtained from American Type Culture Collection (ATCC, Rockville, MD). CHA is an epithelial kidney carcinoma cell line kindly provided by C. Bain (Centre Léon Bérard, Lyon, France). All cell lines were stimulated by PMA-ionomycin for 1 h and 6 h and pooled. Murine fibroblasts transfected with human CD40 ligand (CD40L L cells) were produced in the laboratory (Garrone, et al. (1995) J. Exp. Med. 182:1265-1273). All cell types were cultured in RPMI 1640 (GIBCO BRL, Gaithersburg, MD) supplemented with 10% (vol/vol) heat-inactivated fetal bovine serum (FBS; Flow laboratories, Irvine, UK); 10 mM Hepes, 2 mM L-glutamine, 5 x 10<sup>-5</sup> M 2-mercaptoethanol, penicillin (100 U/ml) and streptomycin (100 µg/ml; hereafter referred to as complete medium).

Generation of DC from CD34<sup>+</sup> cells and from monocytes.

Umbilical cord blood samples were obtained according to appropriate institutional guidelines. Isolation of CD34<sup>+</sup> progenitors was achieved using Minimacs separation columns (Miltenyi Biotec GmbH) as described by Caux, et al. (1996) J. Exp. Med. 184:695-706. In all experiments, the isolated cells were 80 to 99% CD34<sup>+</sup> as judged by staining with anti-CD34 mAb. Cultures of CD34<sup>+</sup> cells

were established in the presence of SCF, GM-CSF, and TNF $\alpha$  as described by Caux, et al. (1992) Nature 360:258-261; or Caux, et al. (1996) J. Exp. Med. 184:695-706. Cells collected after 6 days of culture were separated according to CD1a and CD14 expression into CD14<sup>+</sup>CD1a<sup>-</sup> and CD14<sup>-</sup>CD1a<sup>+</sup> using a FACStar<sup>+</sup> (Becton Dickinson, Mountain View, CA) as described (Caux, et al. (1996) J. Exp. Med. 184:695-706). Cells were further cultured in presence of GM-CSF and TNF $\alpha$  until day 12-17, when 70-90% of cells are CD1a<sup>+</sup> DC. Monocytes were purified by immunomagnetic depletion (Dynal) after preparation of PBMC followed by a 52% Percoll gradient. The depletion was performed with anti-CD3, anti-CD19, and anti-CD8 ascites, and with purified anti-NKH1 (Coulter) and anti-CD16 (Immunotech) mAbs. Monocyte-derived dendritic cells were produced by culturing purified monocytes for 6 days in the presence of GM-CSF and IL-4 (Sallusto and Lanzavecchia (1994) J. Exp. Med. 179:1109-1118). Cells were activated with LPS at the concentration of 25 ng/ml for 1 h to 72 h or with CD40L transfected L cells (one L cell for five DC) (Caux, et al. (1994) J. Exp. Med. 180:1263-1272).

### III. cDNA libraries and isolation of F06B09 cDNA clone

Total RNA was isolated from PMA-ionomycin activated CD1a<sup>-</sup>CD14<sup>+</sup> DC (at day 12 of the culture) and from activated CHA cell line. See Chomczynski and Sacchi (1987) Anal. Biochem. 162:156-159. RNA was treated with DNase I before mRNA purification using the Oligotex-dT kit (Qiagen GmbH, Hilden, Germany). PolyA<sup>+</sup> RNA (2  $\mu$ g) was used to make a cDNA library in the pSport vector (Superscript Plasmid System Kit, GIBCO BRL). A subtraction library was made using the method of Hara et al. (1994) Blood 84:189-199, with minor modifications. In this protocol, tracer (subtracted) cDNA was the CD14<sup>+</sup>-derived DC cDNA, and driver (subtractive) cDNA was CHA cDNA. A 0.6 Kb cDNA containing a polyA tail was isolated from the CD14<sup>+</sup>-derived DC subtraction library. The cDNA of this gene was first amplified using the RACE MARATHON<sup>TM</sup> kit (Clontech, Palo Alto, CA) and two

oligonucleotides: 5'CAGAAATGCCACGAAACAGCCAGGTACT (NGSP1;  
SEQ ID NO: 5) and 5'GCCCCAGTTGCTCATACAAACAGATCAG (GSP1;  
SEQ ID NO: 6) with a recommended cycling program 1. PCR  
products were cloned in the pCRII plasmid (Invitrogen,  
5 San Diego, CA). A lambda CD34<sup>+</sup>-derived DC cDNA library  
was constructed using the GREAT LENGTHS<sup>TM</sup> cDNA Synthesis  
kit with the λTriplex vector (Clontech), and was next  
screened with a 5' F06B09 probe to obtain a full-length  
cDNA. Sequencing was performed on both strands by the  
10 dideoxynucleotide method using a Taq Dye Deoxy Terminator  
Cycle Sequencing kit (Applied Biosystems, Foster City,  
CA) and an automated sequencer (Applied Biosystems).

A clone encoding the human F06B09 protein is  
isolated from a natural human dendritic cell or other  
15 source, by one of many different possible methods. Given  
the sequences provided herein, PCR primers or  
hybridization probes are selected and/or constructed to  
isolate a nucleic acid, e.g., genomic DNA segments or  
cDNA reverse transcripts. Appropriate cell sources  
20 include human tissues, e.g., dendritic cell libraries.  
Tissue distribution below also suggests source tissues.  
Genetic and polymorphic or allelic variants are isolated  
by screening a population of individuals.

This clone was discovered via EST analysis of human  
25 dendritic cell subtraction cDNA library. The driver was  
CD34<sup>+</sup> derived, CD14<sup>+</sup>, PMA, and ionomycin activated  
dendritic cell cDNA, while subtractor was PMA and  
ionomycin activated, kidney carcinoma cell line CHA. The  
initial poly-A containing EST was selected for its  
30 restricted distribution after Northern blot and semi-  
quantitative PCR analysis.

A novel 0.6 Kb partial cDNA (F06B09) was isolated by  
screening a PMA-ionomycin activated CD34<sup>+</sup>-derived DC  
subtraction library. Northern blots probed with the  
35 F06B09 clone showed a 3.7 Kb mRNA transcript  
predominantly expressed in dendritic cells. This mRNA  
was absent by RT-PCR in CHA, the driver epithelial cell  
line used for subtraction. The 5' end of the original  
sequence was extended by RACE and by the screening of a

lambda DC cDNA library, to a final 3691 bp cDNA. This cDNA contains a methionine codon located in a consensus Kozak sequence. See Kozak (1986) Cell. 44:283-292. The full-length cDNA (see Table 1) shows a 342 bp 5'

5 untranslated sequence (nt 1-342), a 1689 bp open reading frame (nt 343-2031), a 3' untranslated sequence of 1660 bp (nt 2032-3691) and a polyadenylation signal AATAAA at position 3638-3643 followed by a poly(A)tail. The encoded protein of 562 amino acids reveals a strong

10 homology with membrane-type matrix metalloproteinase (MT-MMP). As a member of the metalloproteinase family, F06B09 contains a propeptide domain with a cysteine-switch activation domain at Cys69 (Van Wart and Birkedal-Hansen (1990) Proc. Natl. Acad. Sci. USA. 87:5578-5582)

15 and the core enzyme domain contains three zinc-chelating histidine (H) residues at positions 212, 216 and 222 in the zinc binding motif. Like other MT-MMP members, F06B09 presents a consensus insertion RRRR between residues 82-86, corresponding to a furin cleavage site

20 (Table 1). The sequence is followed by a hinge region (260-290) and a potential transmembrane domain of 12 amino acids (526-577) in a hemopexin-like domain (291-538) and a short intracytoplasmic domain (538-541). Multiple alignment with members of the membrane-type

25 matrix metalloproteinase (MT-MMP) family revealed the closest homology with the MT4-MMP (48%) (Puente, et al. (1996) Cancer Res. 56:944-949), and 38%, 39% and 35% respectively with the MT1-MMP (Sato, et al. (1994) Nature 370:61-65) the MT2-MMP (Will and Hinzmann (1995) Eur. J. Biochem. 231:602-608) and the MT3-MMP (Takino, et al. (1995) J. Biol. Chem. 270:23013-23020). Comparison of the most conserved domain, the catalytic domain, showed that F06B09 presents the highest homology to MT4-MMP (48%) (Puente, et al. (1996) Cancer Res. 56:944-949) and

30 significant homologies to other members of the matrix membrane metalloproteinase family (MMP), like the type IV collagenases MMP-9 and MMP-2 (Collier, et al. (1988) J. Biol. Chem. 263:6579-6587; Wilhelm, et al. (1989) J. Biol. Chem. 264:17213-17221).

Therefore, it is suggested that the F06B09 is a fifth member of the MT-MMP subgroup of the MMP.

The coding sequence appears to be complete, encoding a 21 amino acid putative signal peptide followed by a 541  
5 residue polypeptide with significant homology to the membrane-type matrix metalloproteases MT-MMP1 to 4. No evidence yet suggests alternative splicing of this message. The limited EST distribution is indicative of a restricted expression pattern.

10 Further clones will be isolated, e.g., using an antibody based selection procedure. Standard expression cloning methods are applied including, e.g., FACS staining of membrane associated expression product. The antibodies are used to identify clones producing a  
15 recognized protein. Alternatively, antibodies are used to purify an F06B09 protein, with protein sequencing and standard means to isolate a gene encoding that protein.

Genomic or cDNA sequence based methods will also allow for identification of sequences naturally  
20 available, or otherwise, which exhibit homology to the provided sequences.

#### IV. Isolation of mouse F06B09

Similar methods are used as above to isolate an  
25 appropriate F06B09 protein gene. See, e.g., GenBank Accession numbers X91785, X83537, D63579, and U54984. Similar source materials as indicated above are used to isolate natural genes, including genetic, polymorphic, allelic, or strain variants. Species variants are also  
30 isolated using similar methods. Various sequence databases may suggest related or counterpart sequences. See, e.g., Capone, et al. (1996) J. Immunol. 157:969-973.

#### V. Isolation of an avian F06B09 protein clone

35 An appropriate avian source is selected as above. Similar methods are utilized to isolate other species variants, though the level of similarity will typically be lower for avian F06B09 protein as compared to a human to mouse sequence.



## VI. Message distribution

PCR based detection is performed by standard methods, preferably using appropriate primers from opposite ends of the coding sequence, but flanking segments might be selected for specific purposes.

Alternatively, hybridization probes are selected. Particular AT or GC contents of probes are selected depending upon the expected homology and mismatching expected. Appropriate stringency conditions are selected to balance an appropriate positive signal to background ratio. Successive washing steps are used to identify clones of greater homology.

Total RNA (20 µg), extracted from cell lines or cell preparations as described above, were fractionated by electrophoresis on a 1% agarose-formaldehyde gel and transferred onto positively charged nylon membrane (GeneScreenPlus, NEN Life Science Products, Boston, MA) as described by Thomas (1980) Proc. Natl. Acad. Sci. USA. 77:5201-5205. After transfer, blots were cross-linked by UV light (Stratalinker, La Jolla, CA). The original cloned 600 bp fragment was labeled by random priming with <sup>32</sup>P-dCTP (3000 Ci/mmol, Amersham; Ready to Go, Pharmacia Biotech, Orsay, France) and unincorporated nucleotides were removed by spin column chromatography (Chromaspin-100, Clontech). Hybridization and washes were performed in stringent conditions (0.1X SSC/0.1% SDS at 65° C). X-ray films (Kodak, Rochester, NY) were exposed for 3 weeks at -80° C with intensifying screens. Multiple tissue normal fetal and adult organs Northern blots (purchased from Clontech) were similarly used according to the manufacturer's recommendations.

For RT-PCR methods, total RNA extracted from 1 to 10 x 10<sup>6</sup> cells (Chomczynski and Sacchi (1987) Anal. Biochem. 162:156-159) were reverse transcribed using random hexamer primers (Pharmacia, Upsalla, Sweden) and the Superscript RNase-H reverse transcriptase (GIBCO BRL). PCR was performed in a 100 µl volume using 5 ng cDNA, 10 µl 10X PCR reaction buffer (Perkin Elmer Cetus, Norwalk,

(CT), 2.5 U of Taq polymerase (Gene Amp PCR reagents kit: Perkin Elmer Cetus) and 200 mM dNTPs and 500 nM of the 5' and 3' amplification primers. The PCR reactions were made in a DNA thermal cycler (Perkin Elmer) for 35 cycles (1 min denaturation at 94° C, 1 min annealing at 60° C, and 2 min elongation at 72° C).  $\beta$  actin RT-PCR was used as positive control for the efficiency of the reaction using sense and antisense primers. Appropriate sense and antisense primers were used to amplify F06B09. See Table 1.

Northern blot showed a single band of about 4.5 kb in non-activated and PMA and ionomycin-activated, CD34+ derived human DC, and a weak signal in in vitro generated granulocytes. No signal was detected in TF1, Jurkat, CHA, or JY cell lines, nor in freshly isolated monocytes, activated T cells, resting and activated PBLs, or B cells. No expression was found in either fetal or adult tissues. PCR distribution analysis showed expression in activated DC and the MRC5 lung fibroblast cell line, as well as very low signal in U937. The original EST was extended by 5' RACE. A lambda DC cDNA library was screened with a 5' probe. An ORF was identified, which showed highest homology with the human MT4-MMP, a recent addition to the membrane type matrix metalloprotease family. Positive signals were also detected in granulocytes.

Northern blot analysis showed a single ~4 Kb transcript predominantly expressed in resting CD34+-derived DC, to a lesser extent in PMA-ionomycin activated DC, and weakly in granulocytes generated in vitro. The expression pattern of this novel gene was also analyzed by RT-PCR, on freshly isolated cells and on various cell lines. Similarly, RT-PCR analysis confirmed the higher level of F06B09 expression in resting and activated CD34+-derived DC, in granulocytes and to a weaker extent in resting PBLs. F06B09 mRNA is also weakly present on the B cell line, JY. No messenger was detected in TF1 (myelo-erythrocytic), CHA (carcinoma), Jurkat (T cell), MRC5 (fibroblastic) or U937 (myelo-

monocytic) cell lines, nor in freshly isolated monocytes, activated T and B cells or activated PBLs. Among normal human tissues tested, a significant band of ~4 Kb was seen in spleen, lymph node, thymus, appendix, PBL and in bone marrow but absent in fetal tissues. An additional 6 Kb band corresponding probably to unspecific expression or to a longer existing form, was also detected in spleen, PBL and bone marrow but absent in lymph node, thymus, and appendix.

10       The cellular distribution of F06B09 was next determined by the extent of hybridization among the gel-fractionated population of cDNA inserts from libraries made from different cell types. Consistent with the above observations, F06B09 is present in both CD34<sup>+</sup>-derived DC and in monocyte-derived DC, but also in effector T cells, including Th1 and Th2 cells, and to a weaker extent in NK cells. F06B09 mRNA expression is down-regulated after PMA-ionomycin activation at once in DC and T cells. No signal was detected in monocytes, B cell lines nor in different fetal tissues.

In conclusion, the novel MT-MMP appears to be mainly transcribed by resting DC and weakly by effector T cells.

F06B09 mRNA is strongly expressed in different types of DC and down-regulated by CD40L activation. Since the original F06B09 clone was identified in a DC library, further characterization was performed by semi-quantitative RT-PCR. The expression of this gene was analyzed during DC differentiation and maturation, either in DC generated in vitro from CD34<sup>+</sup> cord blood progenitors cultured with GM-CSF and TNF $\alpha$  or from monocytes cultured with GM-CSF and IL-4. During the culture of CD34<sup>+</sup> human cord blood progenitors, F06B09 is first detected at day 6 and increases up to day 12. This messenger is down-regulated after triggering final maturation of the DC by 4 days co-culture with hCD40L-transfected L cells. Similarly, while monocytes do not express detectable amount of F06B09 mRNA, a significant expression could be detected after 6 days of culture in the presence of GM-CSF and IL-4. In contrast,

following activation of these monocyte-derived DC through CD40, the level of mRNA decreases rapidly within 3 h to 12 h. A low amount of F06B09 mRNA is also found in 1h PMA-ionomycin activated CD1a<sup>+</sup> and CD14<sup>+</sup> DC subsets, which is down-regulated after 6h PMA-ionomycin activation. Day 12 macrophages generated in vitro express also weakly F06B09, and 6 h PMA-ionomycin activation of these cells is enough to switch off the signal. In contrast, no signal is detected in freshly isolated Langerhans cells, in basal keratinocytes, in freshly isolated and CD40L activated B cells, and in anti-CD3 and anti-CD28 activated T cells.

Taken together, these results confirm that F06B09 mRNA is expressed in different DC subtypes and rapidly down-regulated upon DC maturation.

#### VII. Chromosomal localization.

The full-length cDNA sequence of F06B09 was analyzed against the EMBL nucleotide and EST databases and resulted in identification of a 436 bp EST (W72721), matching exactly with the F06B09 sequence.

Comparison of the full-length cDNA sequence of F06B09 with the EMBL nucleotide and EST databases identifies a 436 bp EST (W72721), corresponding exactly to the F06B09 sequence. This EST is located on chromosome 16p13.3. In contrast, MT1-MMP and MT3-MMP have been previously located on chromosome 14q11-12 and 8q21.3-22.1 respectively (Mattei, et al. (1997) Genomics. 40:168-169.; Mignon, et al. (1995) Genomics. 28:360-361). Of note, the novel F06B09 MT-MMP gene is on the same chromosome than MT2-MMP, but both genes are located on a different loci; MT2-MMP is on chromosome 16q12 (Mattei, et al. (1997) Genomics. 40:168-169; Yasumitsu, et al. (1997) DNA Res. 4:77-79) whereas the novel MT-MMP is on chromosome 16p13.3.

#### VIII. Expression; purification; characterization

With an appropriate clone from above, the coding sequence is inserted into an appropriate expression

vector. This may be in a vector specifically selected for a prokaryote, yeast, insect, or higher vertebrate, e.g., mammalian expression system. Standard methods are applied to produce the gene product, preferably as a soluble secreted molecule, but will, in certain instances, also be made as an intracellular protein. Intracellular proteins typically require cell lysis to recover the protein, and insoluble inclusion bodies are a common starting material for further purification.

With a clone encoding a vertebrate F06B09 protein, recombinant production means are used, although natural forms may be purified from appropriate sources, e.g., expressing cell lines. The protein product is purified by standard methods of protein purification, in certain cases, e.g., coupled with immunoaffinity methods. Immunoaffinity methods are used either as a purification step, as described above, or as a detection assay to determine the partition properties of the protein.

Preferably, the protein is secreted into the medium, and the soluble product is purified from the medium in a soluble form. Standard purification techniques applied to soluble proteins are then applied, with enzyme assays or immunodetection methods useful for following where the protease purifies. Alternatively, as described above, inclusion bodies from prokaryotic expression systems are a useful source of material. Typically, the insoluble protein is solubilized from the inclusion bodies and refolded using standard methods. Purification methods are developed as described above.

In certain embodiments, the protein is made in a eukaryotic cell which glycosylates the protein normally. The purification methods may be affected thereby, as may biological activities. The intact protein can be processed to release the protein domain, probably due to a cleavage event. While recombinant protein appears to be processed, the physiological processes which normally do such in native cells remain to be determined.

The product of the purification method described above is characterized to determine many structural

features. Standard physical methods are applied, e.g., amino acid analysis and protein sequencing. The resulting protein is subjected to CD spectroscopy and other spectroscopic methods, e.g., NMR, ESR, mass spectroscopy, etc. The product is characterized to determine its molecular form and size, e.g., using gel chromatography and similar techniques. Understanding of the chromatographic properties will lead to more gentle or efficient purification methods.

10        Prediction of glycosylation sites may be made, e.g., as reported in Hansen, et al. (1995) Biochem. J. 308:801-813.

#### 15        IX. Preparation of antibodies against vertebrate F06B09 protein

         With protein produced and purified, as above, animals are immunized to produce antibodies. Polyclonal antiserum may be raised using non-purified antigen, though the resulting serum will exhibit higher background levels. Preferably, the antigen is purified using standard protein purification techniques, including, e.g., affinity chromatography using polyclonal serum indicated above. Presence of specific antibodies is detected using defined synthetic peptide fragments.

25        Alternatively, polyclonal serum is raised against a purified antigen, purified as indicated above, or using synthetic peptides. A series of overlapping synthetic peptides which encompass all of the full length sequence, if presented to an animal, will produce serum recognizing most linear epitopes on the protein. Such an antiserum is used to affinity purify protein. This purified protein, in turn, may be used to immunize another animal to produce another antiserum preparation.

35        Standard techniques are used to generate induce monoclonal antibodies to either unpurified antigen, or, preferably, purified antigen.

#### X. Structure activity relationship

Information on the criticality of particular residues is determined using standard procedures and analysis. Standard mutagenesis analysis is performed, e.g., by generating many different variants at determined positions, e.g., at the positions identified above, and evaluating biological activities of the variants. This may be performed to the extent of determining positions which modify activity, or to focus on specific positions to determine the residues which can be substituted to either retain, block, or modulate biological activity.

Alternatively, analysis of natural variants can indicate what positions tolerate natural mutations. This may result from populational analysis of variation among individuals, or across strains or species. Samples from selected individuals are analyzed, e.g., by PCR analysis and sequencing. This allows evaluation of population polymorphisms.

All references cited herein are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference in its entirety for all purposes.

Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only, and the invention is to be limited only by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled.

## WHAT IS CLAIMED IS:

1. A substantially pure or recombinant F06B09 polypeptide exhibiting at least about 80% sequence  
5 identity over a length of at least about 100 amino acid residues to SEQ ID NO: 4.
2. A fusion protein comprising a polypeptide of claim  
1.  
10
3. A binding compound which specifically binds to a polypeptide of claim 1.
4. The binding compound of claim 3 which is an antibody  
15 or antibody fragment.
5. A nucleic acid encoding a polypeptide of claim 1.
6. An expression vector comprising the nucleic acid of  
20 claim 5.
7. A host cell comprising the vector of claim 6.
8. A process for recombinantly producing a polypeptide  
25 comprising culturing the host cell of claim 7 under conditions in which the polypeptide is expressed.



## SEQUENCE LISTING

- SEQ ID NO: 1 is human F06B-09 nucleotide sequence.  
SEQ ID NO: 2 is human F06B09 amino acid sequence.  
5 SEQ ID NO: 3 is second human F6B09 nucleotide sequence.  
SEQ ID NO: 4 is second human F06B09 amino acid sequence.  
SEQ ID NO: 5 is sense primer.  
SEQ ID NO: 6 is antisense primer  
SEQ ID NO: 7 is primate MT4-MMP amino acid sequence.  
10 SEQ ID NO: 8 is primate MT2-MMP amino acid sequence.  
SEQ ID NO: 9 is primate MT1-MMP amino acid sequence.  
SEQ ID NO: 10 is primate MT3-MMP amino acid sequence.
- 15 (1) GENERAL INFORMATION:  
(i) APPLICANT:  
(A) NAME: Schering Corporation  
(B) STREET: 2000 Galloping Hill Road  
(C) CITY: Kenilworth  
20 (D) STATE: New Jersey  
(E) COUNTRY: USA  
(F) POSTAL CODE (ZIP) 07033  
(G) TELEPHONE (908) 298-5056  
(H) TELEFAX: (908) 298-5388  
25 (ii) TITLE OF INVENTION: MAMMALIAN PROTEINASES;  
RELATED REAGENTS AND METHODS  
(iii) NUMBER OF SEQUENCES:10  
30 (iv) COMPUTER READABLE FORM:  
(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: Power Macintosh  
(C) OPERATING SYSTEM: Mac OS 8.1  
35 (D) SOFTWARE: Microsoft Word 6.0.1  
(v) CURRENT APPLICATION DATA:  
(A) APPLICATION NUMBER:  
(B) FILING DATE:  
40 (C) CLASSIFICATION  
(vi) PRIOR APPLICATION DATA:  
(A) APPLICATION NUMBER: U.S. 09/005,263  
(B) FILING DATE: 09-JAN-1998  
45
- (2) INFORMATION FOR SEQ ID NO:1:  
(i) SEQUENCE CHARACTERISTICS:  
50 (A) LENGTH: 3695 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
55 (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 344..2032

## 5 (ix) FEATURE:

(A) NAME/KEY: mat\_peptide

(B) LOCATION: 398..2032

## 10 (ix) FEATURE:

(A) NAME/KEY: misc\_feature

(B) LOCATION: 3458

(D) OTHER INFORMATION: /note= "nucleotide 3458 designated A, may be A or T"

## 15

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

	CATGCAACAT AATCTTGCTC GATTCTAAAG TCAACGGATC CTGCAAAATT CGCGGCCGCG	60
20	TCAACCCATT AGGTCTTGGC CTTGGAATAA AATTGCTTCT CGTCTGATTC CCGGGCCCAC	120
	CCGACCCAGC GCGCAACCC TGGCCCTCCG GGACCCTCCG CTGACTCCAC CGCGCACTTC	180
	CCGGGACCCC CACACACATC CCAGCCCTCC GGCCGATCCC TCCCTACTCG GTGCCGGGTG	240
25	CCCCCCTTTT TTTTCTAGGC CCGGATCTCC TCCCCAGGT CCCCAGGGCG GCCCCAACCA	300
	GGCCCCCTTC AAACCCCGCC GCGGGCCCGG GCTGGGGCGC ACC ATG CGG CTG CGG	355
30		Met Arg Leu Arg -18 -15
	CTC CGG CTT CTG GCG CTG CTG CTT CTG CAT GCT GGC ACC GCC CGC GCG	403
	Leu Arg Leu Leu Ala Leu Leu Leu Leu His Ala Gly Thr Ala Arg Ala	
	-10 -5 1	
35	CGC CCC GAA GCC CTC GGC GCA GGA CTT AGC CTG GGC TGT GAG AAC TGG	451
	Arg Pro Glu Ala Leu Gly Ala Gly Leu Ser Leu Gly Cys Glu Asn Trp	
	5 10 15	
40	CTG ACT CGC TAT GGT TAC CTA CCG CCA CCC GAC CCT GCC CAG GCC CAG	499
	Leu Thr Arg Tyr Gly Tyr Leu Pro Pro Pro Asp Pro Ala Gln Ala Gln	
	20 25 30	
45	CTG CAG AGC CCT GAA AAT TTG CGC GAT GCC ATC AAA GTC ATG CAA AGG	547
	Leu Gln Ser Pro Glu Asn Leu Arg Asp Ala Ile Lys Val Met Gln Arg	
	35 40 45 50	
	TTC GCG GGG CTG CCG GAG ACC GGC CGC ATG GAC CCA GGG ACA GTG GCC	595
50	Phe Ala Gly Leu Pro Glu Thr Gly Arg Met Asp Pro Gly Thr Val Ala	
	55 60 65	
	ACC ATG CGT AAG CCC CGC TGC TCC CTG CCT GAC GTG CTG GGG GTG GCG	643
	Thr Met Arg Lys Pro Arg Cys Ser Leu Pro Asp Val Leu Gly Val Ala	
	70 75 80	
55	GGG CTG GTC AGG CGG CGT CGC CGG TAC GGT CTG AGC GGC AGC GTG TGG	691
	Gly Leu Val Arg Arg Arg Arg Arg Tyr Gly Leu Ser Gly Ser Val Trp	
	85 90 95	
60	GAG AAG CGA ACC GTG ACA TGG AGG GTA CGT TCC TTC CCC CAG AGC TCC	739
	Glu Lys Arg Thr Val Thr Trp Arg Val Arg Ser Phe Pro Gln Ser Ser	
	100 105 110	
	CAG GTG AGC CAG GAG ACC GTG CGG GTC CTC GTG AGC TAT GCC CTG ATG	787

	Gln	Val	Ser	Gln	Glu	Thr	Val	Arg	Val	Leu	Val	Ser	Tyr	Ala	Leu	Met	
	115					120					125					130	
5	GCG	TGG	GGC	ATG	GAG	TCA	GGC	CTC	ACA	TTT	CAT	GAG	GTG	GAT	TCC	CCC	835
	Ala	Trp	Gly	Met	Glu	Ser	Gly	Leu	Thr	Phe	His	Glu	Val	Asp	Ser	Pro	
					135					140					145		
10	CAG	GGC	CAG	GAG	CCC	GAC	ATC	CTC	ATA	GAC	TTT	GCC	CGC	GCC	TTC	CAA	883
	Gln	Gly	Gln	Glu	Pro	Asp	Ile	Leu	Ile	Asp	Phe	Ala	Arg	Ala	Phe	Gln	
					150					155					160		
15	CAG	GAC	AGC	TAC	CCC	TTC	GAC	GGG	TTG	GGG	GGC	ACC	CTA	GCC	CAT	GCC	931
	Gln	Asp	Ser	Tyr	Pro	Phe	Asp	Gly	Leu	Gly	Gly	Thr	Leu	Ala	His	Ala	
					165				170						175		
20	TTC	TTC	CCT	GGG	GAG	CAC	CCC	ATC	TCC	GGG	GAC	ACT	CAC	TTT	GAC	GAT	979
	Phe	Phe	Pro	Gly	Glu	His	Pro	Ile	Ser	Gly	Asp	Thr	His	Phe	Asp	Asp	
					180				185						190		
25	GAG	GAG	ACC	TGG	ACT	TTT	GGG	TCA	AAA	GAC	GGC	GAG	GGG	ACC	GAC	CTG	1027
	Glu	Glu	Thr	Trp	Thr	Phe	Gly	Ser	Lys	Asp	Gly	Glu	Gly	Thr	Asp	Leu	
						200				205						210	
30	TTT	GCC	GTG	GCT	GTC	CAT	GAG	TTT	GGC	CAC	GCC	CTG	GGC	ATG	GGC	CAC	1075
	Phe	Ala	Val	Ala	Val	His	Glu	Phe	Gly	His	Ala	Leu	Gly	Met	Gly	His	
						215				220					225		
35	TCC	TCA	GCC	CCC	GAC	TCC	ATT	ATG	AGG	CCC	TTC	TAC	CAG	GGT	CCG	GTG	1123
	Ser	Ser	Ala	Pro	Asp	Ser	Ile	Met	Arg	Pro	Phe	Tyr	Gln	Gly	Pro	Val	
						230				235					240		
40	GGC	GAC	CCT	GAC	AAG	TAC	CGC	CTG	TCT	CTG	GAT	GAC	CGC	GAT	GGC	CTG	1171
	Gly	Asp	Pro	Asp	Lys	Tyr	Arg	Leu	Ser	Leu	Asp	Asp	Arg	Asp	Gly	Leu	
						245				250					255		
45	CAG	CAA	CTC	TAT	GGG	AAG	GCG	CCC	CAA	ACC	CCA	TAT	GAC	AAG	CCC	ACA	1219
	Gln	Gln	Leu	Tyr	Gly	Lys	Ala	Pro	Gln	Thr	Pro	Tyr	Asp	Lys	Pro	Thr	
						260				265					270		
50	AGG	AAA	CCC	CTG	GCT	CCT	CCG	CCC	CAG	CCC	CCG	GCC	TCG	CCC	ACA	CAC	1267
	Arg	Lys	Pro	Leu	Ala	Pro	Pro	Pro	Gln	Pro	Pro	Ala	Ser	Pro	Thr	His	
						275				280						290	
55	AGC	CCA	TCC	TTC	CCC	ATC	CCT	GAT	CGA	TGT	GAG	GGC	AAT	TTT	GAC	GCC	1315
	Ser	Pro	Ser	Phe	Pro	Ile	Pro	Asp	Arg	Cys	Glu	Gly	Asn	Phe	Asp	Ala	
						295				300					305		
60	ATC	GCC	AAC	ATC	CGA	GGG	GAA	ACT	TTC	TTC	TTC	AAA	GGC	CCC	TGG	TTC	1363
	Ile	Ala	Asn	Ile	Arg	Gly	Glu	Thr	Phe	Phe	Phe	Lys	Gly	Pro	Trp	Phe	
						310				315					320		
65	TGG	CGC	CTC	CAG	CCC	TCC	GGA	CAG	CTG	GTG	TCC	CCG	CGA	CCC	GCA	CGG	1411
	Trp	Arg	Leu	Gln	Pro	Ser	Gly	Gln	Leu	Val	Ser	Pro	Arg	Pro	Ala	Arg	
						325				330					335		
70	CTG	CAC	CGC	TTC	TGG	GAG	GGG	CTG	CCC	GCC	CAG	GTG	AGG	GTG	GTG	CAG	1459
	Leu	His	Arg	Phe	Trp	Glu	Gly	Leu	Pro	Ala	Gln	Val	Arg	Val	Val	Gln	
						340				345					350		
75	GCC	GCC	TAT	GCT	CGG	CAC	CGA	GAC	GGC	CGA	ATC	CTC	CTC	TTT	AGC	GGG	1507
	Ala	Ala	Tyr	Ala	Arg	His	Arg	Asp	Gly	Arg	Ile	Leu	Leu	Phe	Ser	Gly	
						355				360						370	
80	CCC	CAG	TTC	TGG	GTG	TTC	CAG	GAC	CGG	CAG	CTG	GAG	GGC	GGG	GCG	CGG	1555

	Pro	Gln	Phe	Trp	Val	Phe	Gln	Asp	Arg	Gln	Leu	Glu	Gly	Gly	Ala	Arg	
					375					380					385		
5	CCG	CTC	ACG	GAG	CTG	GGG	CTG	CCC	CCG	GGA	GAG	GAG	GTG	GAC	GCC	GTG	1603
	Pro	Leu	Thr	Glu	Leu	Gly	Leu	Pro	Pro	Gly	Glu	Glu	Val	Asp	Ala	Val	
				390				395						400			
10	TTC	TCG	TGG	CCA	CAG	AAC	GGG	AAG	ACC	TAC	CTG	GTC	CGC	GGC	CGG	CAG	1651
	Phe	Ser	Trp	Pro	Gln	Asn	Gly	Lys	Thr	Tyr	Leu	Val	Arg	Gly	Arg	Gln	
			405				410						415				
15	TAC	TGG	CGC	TAC	GAC	GAG	GCG	GCG	GCG	CGC	CCG	GAC	CCC	GGC	TAC	CTT	1699
	Tyr	Trp	Arg	Tyr	Asp	Glu	Ala	Ala	Ala	Arg	Pro	Asp	Pro	Gly	Tyr	Leu	
		420				425					430						
20	CGC	GAC	CTG	AGC	CTC	TGG	GAA	GGC	GCG	CCC	CCC	TCC	CCT	GAC	GAT	GTC	1747
	Arg	Asp	Leu	Ser	Leu	Trp	Glu	Gly	Ala	Pro	Pro	Ser	Pro	Asp	Asp	Val	
	435				440					445					450		
25	ACC	GTC	AGC	AAC	GCA	GGT	GAC	ACC	TAC	TTC	TTC	AAG	GGC	GCC	CAC	TAC	1795
	Thr	Val	Ser	Asn	Ala	Gly	Asp	Thr	Tyr	Phe	Phe	Lys	Gly	Ala	His	Tyr	
				455				460							465		
30	TGG	CGC	TTC	CCC	AAG	AAC	AGC	ATC	AAG	ACC	GAG	CCG	GAC	GCC	CCC	CAG	1843
	Trp	Arg	Phe	Pro	Lys	Asn	Ser	Ile	Lys	Thr	Glu	Pro	Asp	Ala	Pro	Gln	
			470					475					480				
35	CCC	ATG	GGG	CCC	AAC	TGG	CTG	GAC	TGC	CCC	GCC	CCG	AGC	TCT	GGT	CCC	1891
	Pro	Met	Gly	Pro	Asn	Trp	Leu	Asp	Cys	Pro	Ala	Pro	Ser	Ser	Gly	Pro	
		485					490						495				
40	CGC	GCC	CCC	AGG	CCC	CCC	AAA	GGG	ACC	CCC	GTG	TCC	GAA	ACC	TGC	GAT	1939
	Arg	Ala	Pro	Arg	Pro	Pro	Lys	Gly	Thr	Pro	Val	Ser	Glu	Thr	Cys	Asp	
		500					505				510						
45	TGT	CAG	TGC	GAG	CTC	AAC	CAG	GCC	GCA	GGA	CGT	TGG	CCT	GCT	CCC	ATC	1987
	Cys	Gln	Cys	Glu	Leu	Asn	Gln	Ala	Ala	Gly	Arg	Trp	Pro	Ala	Pro	Ile	
	515				520					525					530		
50	CCG	CTG	CTC	CTC	TTG	CCC	CTG	CTG	GTG	GGG	GGT	GTA	GCC	TCC	CGC		2032
	Pro	Leu	Leu	Leu	Leu	Pro	Leu	Leu	Val	Gly	Gly	Val	Ala	Ser	Arg		
					535				540					545			
55	TGATGGGGGG	AGCCATCCAG	ACCGAACAGC	GCCCTCCACG	GCCGAGTCCC	CCGCCGCTGG											2092
	ACCTGGTTCGG	GGGTTGTGAG	GCGCTGCGGA	GGCCCCCTTGT	CTGTTCCAC	GGACGGGGGC											2152
	TCGGGCGCGG	ACTAAGCAGG	GGGGATCTCC	CGCGCAGGGG	CGGCGGCGGC	GGGGACCGGT											2212
60	CGCCTGGCGC	TGGGCTCAGT	CTCCTCAGGG	TCTGAGACCC	CGGCGCTGCC	ACCGGAACCC											2272
	GCCTTCAGGG	GCGCACGCGC	GCTGGGACCA	TGCGTCGGTC	GTCGCCCCCG	TCGTTCCCTC											2332
	CCGGCTGCCG	CCAGGGGGCG	GTCGGACCCC	GCCTCCCGAG	CCCGGGGAGG	GGCGGGGAGG											2392
65	ACAAGGGGCG	GGCCCCGCGC	CTCACCCGGA	GGGACGGCAG	CCCCGGTCGC	GCGCTGGCCC											2452
	CGCAGGACCT	TCCTTTTCCA	GGAAGAGCCA	GCTTTTCTCG	GAGCGCAGTC	CTGGGACTCT											2512
70	CCGCAGCCCC	GCCCCGCCTG	GCCACTGCGT	CTGGCATTC	TGGGTCGTTA	GAGGACAGGC											2572
	CTGACTGCGA	AGCTGTGCCT	TGCCCCCTCTC	CCACCCGCAG	TTTCTCACCC	CGTTCTGCTC											2632
	CCACAAGGCC	CCCCTACAGT	CACTGCCACA	CTGGTGGGGA	CCTGGGACCC	AGACCCGGAA											2692

5  
 10  
 15  
 20  
 25  
 30  
 35

CCAGCCCAGA TATCACCCTT GAGGACCCAT GCGCCACGTC CTGGGTGGTG GAATCAGTGG 2752  
 GTGGAGGGAC GACCCTTGCT CTCCAGGCTG TTAACCTTTT CCGTTGCTCC CCCGCCACCC 2812  
 ACCTCCTCCT CCCCAGGCCA CCCAACTTGG GCACCTCCCT GGGCCCAGAA CTGCCTTCCA 2872  
 TTCAATGGGG AACCTTCTA TCCCCAAGAA CCCCTTCCCT GCTTGACCCC TGGAGAGAAC 2932  
 AGCTTGACTC CCATCAACTC AACGCTGGTG GAAAGACAGG GACCGAACCC TGGCTCAGGC 2992  
 CTGGTCATTG CCTCCTCAGC ACTCCCTCCT GGGAGGCCTT AGCTCTAGAG TGAGGGGTGG 3052  
 GTGGAACCTG GGGGCACCTC GTTCACCCTG TCCCCACTCC CCACAGTTTT AGGATCTAAA 3112  
 TGATTGCCTC TGGAACTATT CTTCTAGACT ATCCCACATC AGAATCACTG GGAAATTTAA 3172  
 GTTTGCAGAT CCCACACTCA CCCTGAATCC TCACTCAGGG TGGGGTCAGG AATCTGCATT 3232  
 TTAAGTAGTC GCGGGGATTG TGGGGGGCAG TAGCTGGCTG TTTCGTGGCA TTTCTGTGGC 3292  
 TCTGCAGTGT TCCTCCACCC CAGGACCAAT ATGTTTCAGG CACACCGATG GCCTGAACCC 3352  
 CATGGGTAGA GTCACCTAGG GGCCACTTCC TAAGTTGCTG TCCAGCCTCA GTGACCCCTT 3412  
 AGTGCTTCCT GGAGCTGAGG CTGTGGGCGG CTGTCCCAGC AACCAAGCGA GGGGTTGCCC 3472  
 CAGTTGCTCA TACAAACAGA TCAGCATGAG GACAGAAGGC AGGAGACTTT GGTCACTTAC 3532  
 CTGGGAATTC TGGGCTGCCA GGAAACGATT TGGGCCTCTG TCAGTTTCTT TTCCATGTAT 3592  
 GAGGAGGGGG AAATTTGTAT ATTAGATACT TATTCATCCC ACTCTGGACA ATAAAAACGA 3652  
 ATGTACAAAA AAAACATAAA AAAAAAAAT AAAGAAAATC AAA 3695

## (2) INFORMATION FOR SEQ ID NO:2:

40  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 563 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

45  
 (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

50  
 55  
 60

Met Arg Leu Arg Leu Arg Leu Leu Ala Leu Leu Leu Leu His Ala Gly  
 -18 -15 -10 -5  
 Thr Ala Arg Ala Arg Pro Glu Ala Leu Gly Ala Gly Leu Ser Leu Gly  
 1 5 10  
 Cys Glu Asn Trp Leu Thr Arg Tyr Gly Tyr Leu Pro Pro Pro Asp Pro  
 15 20 25 30  
 Ala Gln Ala Gln Leu Gln Ser Pro Glu Asn Leu Arg Asp Ala Ile Lys  
 35 40 45  
 Val Met Gln Arg Phe Ala Gly Leu Pro Glu Thr Gly Arg Met Asp Pro  
 50 55 60  
 Gly Thr Val Ala Thr Met Arg Lys Pro Arg Cys Ser Leu Pro Asp Val  
 65 70 75

Leu Gly Val Ala Gly Leu Val Arg Arg Arg Arg Arg Tyr Gly Leu Ser  
 80 85 90

5 Gly Ser Val Trp Glu Lys Arg Thr Val Thr Trp Arg Val Arg Ser Phe  
 95 100 105 110

Pro Gln Ser Ser Gln Val Ser Gln Glu Thr Val Arg Val Leu Val Ser  
 115 120 125

10 Tyr Ala Leu Met Ala Trp Gly Met Glu Ser Gly Leu Thr Phe His Glu  
 130 135 140

Val Asp Ser Pro Gln Gly Gln Glu Pro Asp Ile Leu Ile Asp Phe Ala  
 145 150 155

Arg Ala Phe Gln Gln Asp Ser Tyr Pro Phe Asp Gly Leu Gly Gly Thr  
 160 165 170

20 Leu Ala His Ala Phe Phe Pro Gly Glu His Pro Ile Ser Gly Asp Thr  
 175 180 185 190

His Phe Asp Asp Glu Glu Thr Trp Thr Phe Gly Ser Lys Asp Gly Glu  
 195 200 205

25 Gly Thr Asp Leu Phe Ala Val Ala Val His Glu Phe Gly His Ala Leu  
 210 215 220

Gly Met Gly His Ser Ser Ala Pro Asp Ser Ile Met Arg Pro Phe Tyr  
 225 230 235

30 Gln Gly Pro Val Gly Asp Pro Asp Lys Tyr Arg Leu Ser Leu Asp Asp  
 240 245 250

35 Arg Asp Gly Leu Gln Gln Leu Tyr Gly Lys Ala Pro Gln Thr Pro Tyr  
 255 260 265 270

Asp Lys Pro Thr Arg Lys Pro Leu Ala Pro Pro Pro Gln Pro Pro Ala  
 275 280 285

40 Ser Pro Thr His Ser Pro Ser Phe Pro Ile Pro Asp Arg Cys Glu Gly  
 290 295 300

Asn Phe Asp Ala Ile Ala Asn Ile Arg Gly Glu Thr Phe Phe Phe Lys  
 305 310 315

Gly Pro Trp Phe Trp Arg Leu Gln Pro Ser Gly Gln Leu Val Ser Pro  
 320 325 330

50 Arg Pro Ala Arg Leu His Arg Phe Trp Glu Gly Leu Pro Ala Gln Val  
 335 340 345 350

Arg Val Val Gln Ala Ala Tyr Ala Arg His Arg Asp Gly Arg Ile Leu  
 355 360 365

55 Leu Phe Ser Gly Pro Gln Phe Trp Val Phe Gln Asp Arg Gln Leu Glu  
 370 375 380

Gly Gly Ala Arg Pro Leu Thr Glu Leu Gly Leu Pro Pro Gly Glu Glu  
 385 390 395

60 Val Asp Ala Val Phe Ser Trp Pro Gln Asn Gly Lys Thr Tyr Leu Val  
 400 405 410

Arg Gly Arg Gln Tyr Trp Arg Tyr Asp Glu Ala Ala Ala Arg Pro Asp  
 415 420 425 430  
 5 Pro Gly Tyr Leu Arg Asp Leu Ser Leu Trp Glu Gly Ala Pro Pro Ser  
 435 440 445  
 Pro Asp Asp Val Thr Val Ser Asn Ala Gly Asp Thr Tyr Phe Phe Lys  
 450 455 460  
 10 Gly Ala His Tyr Trp Arg Phe Pro Lys Asn Ser Ile Lys Thr Glu Pro  
 465 470 475  
 Asp Ala Pro Gln Pro Met Gly Pro Asn Trp Leu Asp Cys Pro Ala Pro  
 480 485 490  
 15 Ser Ser Gly Pro Arg Ala Pro Arg Pro Pro Lys Gly Thr Pro Val Ser  
 495 500 505 510  
 20 Glu Thr Cys Asp Cys Gln Cys Glu Leu Asn Gln Ala Ala Gly Arg Trp  
 515 520 525  
 Pro Ala Pro Ile Pro Leu Leu Leu Leu Pro Leu Leu Val Gly Gly Val  
 530 535 540  
 25 Ala Ser Arg  
 545

## (2) INFORMATION FOR SEQ ID NO:3:

30 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 3691 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: cDNA

40 (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 343..2028

45 (ix) FEATURE:  
 (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 406..2028

50 (ix) FEATURE:  
 (A) NAME/KEY: misc\_feature  
 (B) LOCATION: 3454  
 (D) OTHER INFORMATION: /note= "nucleotide 3454 is W,  
 designated C herein"

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CATGCAACAT AATCTTGCTC GATTCTAAAG TCAACGGATC CTGCAAAATT CGCGGCCGCG	60
TCAACCCATT AGGTCTTGGC CTTGGAATAA AATTGCTTCT CGTCTGATTC CCGGGCCCAC	120
60 CCGACCCAGC GGCGCAACCC TGGCCCTCCG GGACCCTCCG CTGACTCCAC CGCGCACTTC	180
CCGGGACCCC CACACACATC CCAGCCCTCC GGCCGATCCC TCCCTACTCG GTGCCGGGTG	240
CCCCCGCCCC TCTCCAGGCC CGGATCTCCT CCCCAGGTC CCCGGGGCGG CCCCAGCCAG	300

	GCCCCCTTCG AACCCCGCCG GCGGCCCGGG CTGGGGCGCA CC ATG CGG CTG CGG	354
	Met Arg Leu Arg	
	-21 -20	
5	CTC CGG CTT CTG GCG CTG CTG CTT CTG CTG CTG GCA CCG CCC GCG CGC	402
	Leu Arg Leu Leu Ala Leu Leu Leu Leu Leu Ala Pro Pro Ala Arg	
	-15 -10 -5	
10	GCC CCG AAG CCC TCG GCG CAG GAC GTG AGC CTG GGC GTG GAC TGG CTG	450
	Ala Pro Lys Pro Ser Ala Gln Asp Val Ser Leu Gly Val Asp Trp Leu	
	1 5 10 15	
15	ACT CGC TAT GGT TAC CTG CCG CCA CCC CAC CCT GCC CAG GCC CAG CTG	498
	Thr Arg Tyr Gly Tyr Leu Pro Pro Pro His Pro Ala Gln Ala Gln Leu	
	20 25 30	
20	CAG AGC CCT GAG AAG TTG CGC GAT GCC ATC AAA GTC ATG CAG AGG TTC	546
	Gln Ser Pro Glu Lys Leu Arg Asp Ala Ile Lys Val Met Gln Arg Phe	
	35 40 45	
25	GCG GGG CTG CCG GAG ACC GGC CGC ATG GAC CCA GGG ACA GTG GCC ACC	594
	Ala Gly Leu Pro Glu Thr Gly Arg Met Asp Pro Gly Thr Val Ala Thr	
	50 55 60	
30	ATG CGT AAG CCC CGC TGC TCC CTG CCT GAC GTG CTG GGG GTG GCG GGG	642
	Met Arg Lys Pro Arg Cys Ser Leu Pro Asp Val Leu Gly Val Ala Gly	
	65 70 75	
35	CTG GTC AGG CGG CGT CGC CGG TAC GCT CTG AGC GGC AGC GTG TGG AAG	690
	Leu Val Arg Arg Arg Arg Arg Tyr Ala Leu Ser Gly Ser Val Trp Lys	
	80 85 90 95	
40	AAG CGA ACC CTG ACA TGG AGG GTA CGT TCC TTC CCC CAG AGC TCC CAG	738
	Lys Arg Thr Leu Thr Trp Arg Val Arg Ser Phe Pro Gln Ser Ser Gln	
	100 105 110	
45	CTG AGC CAG GAG ACC GTG CGG GTC CTC ATG AGC TAT GCC CTG ATG GCC	786
	Leu Ser Gln Glu Thr Val Arg Val Leu Met Ser Tyr Ala Leu Met Ala	
	115 120 125	
50	TGG GGC ATG GAG TCA GGC CTC ACA TTT CAT GAG GTG GAT TCC CCC CAG	834
	Trp Gly Met Glu Ser Gly Leu Thr Phe His Glu Val Asp Ser Pro Gln	
	130 135 140	
55	GGC CAG GAG CCC GAC ATC CTC ATC GAC TTT GCC CGC GCC TTC CAC CAG	882
	Gly Gln Glu Pro Asp Ile Leu Ile Asp Phe Ala Arg Ala Phe His Gln	
	145 150 155	
60	GAC AGC TAC CCC TTC GAC GGG TTG GGG GGC ACC CTA GCC CAT GCC TTC	930
	Asp Ser Tyr Pro Phe Asp Gly Leu Gly Gly Thr Leu Ala His Ala Phe	
	160 165 170 175	
65	TTC CCT GGG GAG CAC CCC ATC TCC GGG GAC ACT CAC TTT GAC GAT GAG	978
	Phe Pro Gly Glu His Pro Ile Ser Gly Asp Thr His Phe Asp Asp Glu	
	180 185 190	
70	GAG ACC TGG ACT TTT GGG TCA AAA GAC GGC GAG GGG ACC GAC CTG TTT	1026
	Glu Thr Trp Thr Phe Gly Ser Lys Asp Gly Glu Gly Thr Asp Leu Phe	
	195 200 205	
75	GCC GTG GCT GTC CAT GAG TTT GGC CAC GCC CTG GGC CTG GGC CAC TTC	1074
	Ala Val Ala Val His Glu Phe Gly His Ala Leu Gly Leu Gly His Ser	
	210 215 220	



5	TC	A	GCC	CCC	AAC	TCC	ATT	ATG	AGG	CCC	TTC	TAC	CAG	GGT	CCG	GTG	GGC	1122
	Ser	Ala	Pro	Asn	Ser	Ile	Met	Arg	Pro	Phe	Tyr	Gln	Gly	Pro	Val	Gly		
	225						230					235						
10	GAC	CCT	GAC	AAG	TAC	CGC	CTG	TCT	CAG	GAT	GAC	CGC	GAT	GGC	CTG	CAG	1170	
	Asp	Pro	Asp	Lys	Tyr	Arg	Leu	Ser	Gln	Asp	Asp	Arg	Asp	Gly	Leu	Gln		
	240					245				250					255			
15	CAA	CTC	TAT	GGG	AAG	GCG	CCC	CAA	ACC	CCA	TAT	GAC	AAG	CCC	ACA	AGG	1218	
	Gln	Leu	Tyr	Gly	Lys	Ala	Pro	Gln	Thr	Pro	Tyr	Asp	Lys	Pro	Thr	Arg		
				260						265					270			
20	AAA	CCC	CTG	GCT	CCT	CCG	CCC	CAG	CCC	CCG	GCC	TCG	CCC	ACA	CAC	AGC	1266	
	Lys	Pro	Leu	Ala	Pro	Pro	Pro	Gln	Pro	Pro	Ala	Ser	Pro	Thr	His	Ser		
			275					280						285				
25	CCA	TCC	TTC	CCC	ATC	CCT	GAT	CGA	TGT	GAG	GGC	AAT	TTT	GAC	GCC	ATC	1314	
	Pro	Ser	Phe	Pro	Ile	Pro	Asp	Arg	Cys	Glu	Gly	Asn	Phe	Asp	Ala	Ile		
			290					295					300					
30	GCC	AAC	ATC	CGA	GGG	GAA	ACT	TTC	TTC	TTC	AAA	GGC	CCC	TGG	TTC	TGG	1362	
	Ala	Asn	Ile	Arg	Gly	Glu	Thr	Phe	Phe	Phe	Lys	Gly	Pro	Trp	Phe	Trp		
		305					310					315						
35	CGC	CTC	CAG	CCC	TCC	GGA	CAG	CTG	GTG	TCC	CCG	CGA	CCC	GCA	CGG	CTG	1410	
	Arg	Leu	Gln	Pro	Ser	Gly	Gln	Leu	Val	Ser	Pro	Arg	Pro	Ala	Arg	Leu		
	320					325					330					335		
40	CAC	CGC	TTC	TGG	GAG	GGG	CTG	CCC	GCC	CAG	GTG	AGG	GTG	GTG	CAG	GCC	1458	
	His	Arg	Phe	Trp	Glu	Gly	Leu	Pro	Ala	Gln	Val	Arg	Val	Val	Gln	Ala		
					340					345					350			
45	GCC	TAT	GCT	CGG	CAC	CGA	GAC	GGC	CGA	ATC	CTC	CTC	TTT	AGC	GGG	CCC	1506	
	Ala	Tyr	Ala	Arg	His	Arg	Asp	Gly	Arg	Ile	Leu	Leu	Phe	Ser	Gly	Pro		
			355					360						365				
50	CAG	TTC	TGG	GTG	TTC	CAG	GAC	CGG	CAG	CTG	GAG	GGC	GGG	GCG	CGG	CCG	1554	
	Gln	Phe	Trp	Val	Phe	Gln	Asp	Arg	Gln	Leu	Glu	Gly	Gly	Ala	Arg	Pro		
			370					375					380					
55	CTC	ACG	GAG	CTG	GGG	CTG	CCC	CCG	GGA	GAG	GAG	GTG	GAC	GCC	GTG	TTC	1602	
	Leu	Thr	Glu	Leu	Gly	Leu	Pro	Pro	Gly	Glu	Glu	Val	Asp	Ala	Val	Phe		
		385					390					395						
60	TCG	TGG	CCA	CAG	AAC	GGG	AAG	ACC	TAC	CTG	GTC	CGC	GGC	CGG	CAG	TAC	1650	
	Ser	Trp	Pro	Gln	Asn	Gly	Lys	Thr	Tyr	Leu	Val	Arg	Gly	Arg	Gln	Tyr		
	400					405					410					415		
65	TGG	CGC	TAC	GAC	GAG	GCG	GCG	GCG	CGC	CCG	GAC	CCC	GGC	TAC	CCT	CGC	1698	
	Trp	Arg	Tyr	Asp	Glu	Ala	Ala	Ala	Arg	Pro	Asp	Pro	Gly	Tyr	Pro	Arg		
					420					425					430			
70	GAC	CTG	AGC	CTC	TGG	GAA	GGC	GCG	CCC	CCC	TCC	CCT	GAC	GAT	GTC	ACC	1746	
	Asp	Leu	Ser	Leu	Trp	Glu	Gly	Ala	Pro	Pro	Ser	Pro	Asp	Asp	Val	Thr		
				435					440					445				
75	GTC	AGC	AAC	GCA	GGT	GAC	ACC	TAC	TTC	TTC	AAG	GGC	GCC	CAC	TAC	TGG	1794	
	Val	Ser	Asn	Ala	Gly	Asp	Thr	Tyr	Phe	Phe	Lys	Gly	Ala	His	Tyr	Trp		
			450					455					460					
80	CGC	TTC	CCC	AAG	AAC	AGC	ATC	AAG	ACC	GAG	CCG	GAC	GCC	CCC	CAG	CCC	1842	
	Arg	Phe	Pro	Lys	Asn	Ser	Ile	Lys	Thr	Glu	Pro	Asp	Ala	Pro	Gln	Pro		
		465					470					475						

	ATG GGG CCC AAC TGG CTG GAC TGC CCC GCC CCG AGC TCT GGT CCC CGC	1890
	Met Gly Pro Asn Trp Leu Asp Cys Pro Ala Pro Ser Ser Gly Pro Arg	
	480 485 490 495	
5	GCC CCC AGG CCC CCC AAA GCG ACC CCC GTG TCC GAA ACC TGC GAT TGT	1938
	Ala Pro Arg Pro Pro Lys Ala Thr Pro Val Ser Glu Thr Cys Asp Cys	
	500 505 510	
10	CAG TGC GAG CTC AAC CAG GCC GCA GGA CGT TGG CCT GCT CCC ATC CCG	1986
	Gln Cys Glu Leu Asn Gln Ala Ala Gly Arg Trp Pro Ala Pro Ile Pro	
	515 520 525	
15	CTG CTC CTC TTG CCC CTG CTG GTG GGG GGT GTA GCC TCC CGC	2028
	Leu Leu Leu Leu Pro Leu Leu Val Gly Gly Val Ala Ser Arg	
	530 535 540	
	TGATGGGGGG AGCCATCCAG ACCGAACAGC GCCCTCCACG GCCGAGTCCC CCGCCGCTGG	2088
20	ACCTGGTCGG GGGTTGTGAG GCGCTGCGGA GGCCCCCTTGT CTGTTCCAC GGACGGGGGC	2148
	TCGGGCGCGG ACTAAGCAGG GGGGATCTCC CGCGCAGGGG CGGCGGCGGC GGGGACCGGT	2208
	CGCCTGGCGC TGGGCTCAGT CTCCTCAGGG TCTGAGACCC CGGCGCTGCC ACCGGAACCC	2268
25	GCCTTCAGGG GCGCACGCGC GCTGGGACCA TGCCTCGGTC GTCGCCCCCG TCGTTCCCTC	2328
	CCGGCTGCCG CCAGGGGGCG GTCGGACCCC GCCTCCCGAG CCCGGGGAGG GGCGGGGAGG	2388
30	ACAAGGGGCG GGCCCGCGGC CTCACCCGGA GGGACGGCAG CCCCAGTCCG GCGCTGGCCC	2448
	CGCAGGACCT TCCTTTTCCA GGAAGAGCCA GCTTTTCTCG GAGCGCAGTC CTGGGACTCT	2508
	CCGCAGCCCC GCCCCGCCTG GCCACTGCGT CTGGCATTC TGGGTCGTTA GAGGACAGGC	2568
35	CTGACTGCGA AGCTGTGCCT TGCCCCCTCTC CCACCCGCAG TTTCTCACCC CGTTCTGCTC	2628
	CCACAAGGCC CCCCTACAGT CACTGCCACA CTGGTGGGGA CCTGGGACCC AGACCCGGAA	2688
40	CCAGCCCAGA TATCACCCCT GAGGACCCAT GCGCCACGTC CTGGGTGGTG GAATCAGTGG	2748
	CTGGAGGGAC GACCCTTGCT CTCCAGGCTG TTAACCTTTT CCGTTGCTCC CCCGCCACCC	2808
	ACCTCCTCCT CCCAGGCCA CCCAACTTGG GCACCTCCCT GGGCCCAGAA CTGCCTTCCA	2868
45	TTCAATGGGG AACCCTTCTA TCCCCAAGAA CCCCTTCCCT GCTTGACCCC TGGAGAGAAC	2928
	AGCTTGACTC CCATCAACTC AACGCTGGTG GAAAGACAGG GACCGAACCC TGGCTCAGGC	2988
50	CTGGTCATTG CCTCCTCAGC ACTCCCTCCT GGGAGGCCCT AGCTCTAGAG TGAGGGGTGG	3048
	GTGGAACCTG GGGGCACCTC GTTCACCCTG TCCCCACTCC CCACAGTTTT AGGATCTAAA	3108
	TGATTGCCTC TGGAATATT CTTCTAGACT ATCCACATC AGAATCACTG GGAAATTTAA	3168
55	GTTTGCAGAT CCCACACTCA CCCTGAATCC TCACTCAGGG TGGGGTCAGG AATCTGCATT	3228
	TTAACTAGTC GCGGGGATTG TGGGGGGCAG TAGCTGGCTG TTTCTGTGGC TTTCTGTGGC	3288
60	TCTGCAGTGT TCCTCCACCC CAGGACCAAT ATGTTTCAGGC CACACCGATG GCCTGAACCC	3348
	CATGGGTAGA GTCACCTAGG GGCCACTTCC TAAGTTGCTG TCCAGCCTCA GTGACCCCTT	3408
	AGTGCTTCCT GGAGCTGAGG CTGTGGGCGG CTGTCCCAGC AACCACGCGA GGGGTTGCC	3468

CAGTTGCTCA TACAAACAGA TCAGCATGAG GACAGAAGGC AGGAGACTTT GGTCAGTTAC 3528  
 CTGGGAATTC TGGGCTGCCA GGAAACGATT TGGGCCTCTG TCAGTTTCTT TTCCATGTAT 3588  
 5 GAGGAGGGGG AAATTTGTAT ATTAGATACT TATTCATCCC ACTCTGGACA ATAAAAACGA 3648  
 ATGTACAAAA AAAACATAAA AAAAAAAAT AAAGAAAATC AAA 3691

10

(2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

15

- (A) LENGTH: 562 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

20

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Arg Leu Arg Leu Arg Leu Leu Ala Leu Leu Leu Leu Leu Ala  
 -21 -20 -15 -10  
 25 Pro Pro Ala Arg Ala Pro Lys Pro Ser Ala Gln Asp Val Ser Leu Gly  
 -5 1 5 10  
 Val Asp Trp Leu Thr Arg Tyr Gly Tyr Leu Pro Pro Pro His Pro Ala  
 15 20 25  
 30 Gln Ala Gln Leu Gln Ser Pro Glu Lys Leu Arg Asp Ala Ile Lys Val  
 30 35 40  
 35 Met Gln Arg Phe Ala Gly Leu Pro Glu Thr Gly Arg Met Asp Pro Gly  
 45 50 55  
 Thr Val Ala Thr Met Arg Lys Pro Arg Cys Ser Leu Pro Asp Val Leu  
 60 65 70 75  
 40 Gly Val Ala Gly Leu Val Arg Arg Arg Arg Arg Tyr Ala Leu Ser Gly  
 80 85 90  
 Ser Val Trp Lys Lys Arg Thr Leu Thr Trp Arg Val Arg Ser Phe Pro  
 95 100 105  
 45 Gln Ser Ser Gln Leu Ser Gln Glu Thr Val Arg Val Leu Met Ser Tyr  
 110 115 120  
 50 Ala Leu Met Ala Trp Gly Met Glu Ser Gly Leu Thr Phe His Glu Val  
 125 130 135  
 Asp Ser Pro Gln Gly Gln Glu Pro Asp Ile Leu Ile Asp Phe Ala Arg  
 140 145 150 155  
 55 Ala Phe His Gln Asp Ser Tyr Pro Phe Asp Gly Leu Gly Gly Thr Leu  
 160 165 170  
 Ala His Ala Phe Phe Pro Gly Glu His Pro Ile Ser Gly Asp Thr His  
 175 180 185  
 60 Phe Asp Asp Glu Glu Thr Trp Thr Phe Gly Ser Lys Asp Gly Glu Gly  
 190 195 200  
 Thr Asp Leu Phe Ala Val Ala Val His Glu Phe Gly His Ala Leu Gly

	205	210	215
	Leu Gly His Ser Ser Ala Pro Asn Ser Ile Met Arg Pro Phe Tyr Gln		
	220	225	230 235
5	Gly Pro Val Gly Asp Pro Asp Lys Tyr Arg Leu Ser Gln Asp Asp Arg		
		240 245	250
10	Asp Gly Leu Gln Gln Leu Tyr Gly Lys Ala Pro Gln Thr Pro Tyr Asp		
		255 260	265
	Lys Pro Thr Arg Lys Pro Leu Ala Pro Pro Pro Gln Pro Pro Ala Ser		
		270 275	280
15	Pro Thr His Ser Pro Ser Phe Pro Ile Pro Asp Arg Cys Glu Gly Asn		
		285 290	295
	Phe Asp Ala Ile Ala Asn Ile Arg Gly Glu Thr Phe Phe Phe Lys Gly		
	300	305	310 315
20	Pro Trp Phe Trp Arg Leu Gln Pro Ser Gly Gln Leu Val Ser Pro Arg		
		320 325	330
	Pro Ala Arg Leu His Arg Phe Trp Glu Gly Leu Pro Ala Gln Val Arg		
25		335 340	345
	Val Val Gln Ala Ala Tyr Ala Arg His Arg Asp Gly Arg Ile Leu Leu		
		350 355	360
30	Phe Ser Gly Pro Gln Phe Trp Val Phe Gln Asp Arg Gln Leu Glu Gly		
		365 370	375
	Gly Ala Arg Pro Leu Thr Glu Leu Gly Leu Pro Pro Gly Glu Glu Val		
	380	385 390	395
35	Asp Ala Val Phe Ser Trp Pro Gln Asn Gly Lys Thr Tyr Leu Val Arg		
		400 405	410
	Gly Arg Gln Tyr Trp Arg Tyr Asp Glu Ala Ala Ala Arg Pro Asp Pro		
40		415 420	425
	Gly Tyr Pro Arg Asp Leu Ser Leu Trp Glu Gly Ala Pro Pro Ser Pro		
		430 435	440
45	Asp Asp Val Thr Val Ser Asn Ala Gly Asp Thr Tyr Phe Phe Lys Gly		
		445 450	455
	Ala His Tyr Trp Arg Phe Pro Lys Asn Ser Ile Lys Thr Glu Pro Asp		
	460	465 470	475
50	Ala Pro Gln Pro Met Gly Pro Asn Trp Leu Asp Cys Pro Ala Pro Ser		
		480 485	490
	Ser Gly Pro Arg Ala Pro Arg Pro Pro Lys Ala Thr Pro Val Ser Glu		
55		495 500	505
	Thr Cys Asp Cys Gln Cys Glu Leu Asn Gln Ala Ala Gly Arg Trp Pro		
		510 515	520
60	Ala Pro Ile Pro Leu Leu Leu Leu Pro Leu Leu Val Gly Gly Val Ala		
		525 530	535
	Ser Arg		
	540		

## (2) INFORMATION FOR SEQ ID NO:5:

- 5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- 10 (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "primer"

15

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CAGAAATGCC ACGAAACAGC CAGGTACT

28

## 20 (2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28 base pairs  
(B) TYPE: nucleic acid  
25 (C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "primer"

30

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

35

GCCCCAGTTG CTCATACAAA CAGATCAG

28

## (2) INFORMATION FOR SEQ ID NO:7:

- 40 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 519 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: not relevant  
(D) TOPOLOGY: linear

45

## (ii) MOLECULE TYPE: peptide

50

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met Gln Gln Phe Gly Gly Leu Glu Ala Thr Gly Ile Leu Asp Glu Ala  
1 5 10 15

Thr Leu Ala Leu Met Lys Thr Pro Arg Cys Ser Leu Pro Asp Leu Pro  
20 25 30

Val Leu Thr Gln Ala Arg Arg Arg Arg Gln Ala Pro Ala Pro Thr Lys  
35 40 45

Trp Asn Lys Arg Asn Leu Ser Trp Arg Val Arg Thr Phe Pro Arg Asp  
50 55 60

	Ser	Pro	Leu	Gly	His	Asp	Thr	Val	Arg	Ala	Leu	Met	Tyr	Tyr	Ala	Leu	
	65					70					75					80	
5	Lys	Val	Trp	Ser	Asp	Ile	Ala	Pro	Leu	Asn	Phe	His	Glu	Val	Ala	Gly	
					85					90					95		
	Ser	Thr	Ala	Asp	Ile	Gln	Ile	Asp	Phe	Ser	Lys	Ala	Asp	His	Asn	Asp	
				100					105					110			
10	Gly	Tyr	Pro	Phe	Asp	Gly	Pro	Gly	Gly	Thr	Val	Ala	His	Ala	Phe	Phe	
			115					120					125				
	Pro	Gly	His	His	His	Thr	Ala	Gly	Asp	Thr	His	Phe	Asp	Asp	Asp	Glu	
15			130				135					140					
	Ala	Trp	Thr	Phe	Arg	Ser	Ser	Asp	Ala	His	Gly	Met	Asp	Leu	Phe	Ala	
	145					150					155					160	
20	Val	Ala	Val	His	Glu	Phe	Gly	His	Ala	Ile	Gly	Leu	Ser	His	Val	Ala	
				165						170					175		
	Ala	Ala	His	Ser	Ile	Met	Arg	Pro	Tyr	Tyr	Gln	Gly	Pro	Val	Gly	Asp	
				180					185					190			
25	Pro	Leu	Arg	Tyr	Gly	Leu	Pro	Tyr	Glu	Asp	Lys	Val	Arg	Val	Trp	Gln	
			195					200					205				
	Leu	Tyr	Gly	Val	Arg	Glu	Ser	Val	Ser	Pro	Thr	Ala	Gln	Pro	Glu	Glu	
30		210					215					220					
	Pro	Pro	Leu	Leu	Pro	Glu	Pro	Pro	Asp	Asn	Arg	Ser	Ser	Ala	Pro	Pro	
	225					230				235						240	
35	Arg	Lys	Asp	Val	Pro	His	Arg	Cys	Ser	Thr	His	Phe	Asp	Ala	Val	Ala	
				245						250					255		
	Gln	Ile	Arg	Gly	Glu	Ala	Phe	Phe	Phe	Lys	Gly	Lys	Tyr	Phe	Trp	Arg	
				260					265					270			
40	Leu	Thr	Arg	Asp	Arg	His	Leu	Val	Ser	Leu	Gln	Pro	Ala	Gln	Met	His	
			275					280					285				
	Arg	Phe	Trp	Arg	Gly	Leu	Pro	Leu	His	Leu	Asp	Ser	Val	Asp	Ala	Val	
45		290					295					300					
	Tyr	Glu	Arg	Thr	Ser	Asp	His	Lys	Ile	Val	Phe	Phe	Lys	Gly	Asp	Arg	
	305					310					315					320	
50	Tyr	Trp	Val	Phe	Lys	Asp	Asn	Asn	Val	Glu	Glu	Gly	Tyr	Pro	Arg	Pro	
				325						330				335			
	Val	Ser	Asp	Phe	Ser	Leu	Pro	Pro	Gly	Gly	Ile	Asp	Ala	Ala	Phe	Ser	
				340					345					350			
55	Trp	Ala	His	Asn	Asp	Arg	Thr	Tyr	Phe	Phe	Lys	Asp	Gln	Leu	Tyr	Trp	
			355					360					365				
	Arg	Tyr	Asp	Asp	His	Thr	Arg	His	Met	Asp	Pro	Gly	Tyr	Pro	Ala	Gln	
60		370					375					380					
	Ser	Pro	Leu	Trp	Arg	Gly	Val	Pro	Ser	Thr	Leu	Asp	Asp	Ala	Met	Arg	
	385					390					395					400	
	Trp	Ser	Asp	Gly	Ala	Ser	Tyr	Phe	Phe	Arg	Gly	Gln	Glu	Tyr	Trp	Lys	

405 410 415

Val Leu Asp Gly Glu Leu Glu Val Ala Pro Gly Tyr Pro Gln Ser Thr  
420 425 430

5 Ala Arg Asp Trp Leu Val Cys Gly Asp Ser Gln Ala Asp Gly Ser Val  
435 440 445

10 Ala Ala Gly Val Asp Ala Ala Glu Gly Pro Arg Ala Pro Pro Gly Gln  
450 455 460

His Asp Gln Ser Arg Ser Glu Asp Gly Tyr Glu Val Cys Ser Cys Thr  
465 470 475 480

15 Ser Gly Ala Ser Ser Pro Pro Gly Ala Pro Gly Pro Leu Val Ala Ala  
485 490 495

Thr Met Leu Leu Leu Leu Pro Pro Leu Ser Pro Gly Ala Leu Trp Thr  
500 505 510

20 Ala Ala Gln Ala Leu Thr Leu  
515

## (2) INFORMATION FOR SEQ ID NO:8:

25

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 564 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: not relevant

30

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

35

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

40 Met Lys Arg Pro Arg Cys Gly Val Pro Asp Gln Phe Gly Val Arg Val  
1 5 10 15

Lys Ala Asn Leu Arg Arg Arg Arg Lys Arg Tyr Ala Leu Thr Gly Arg  
20 25 30

45 Lys Trp Asn Asn His His Leu Thr Phe Ser Ile Gln Asn Tyr Thr Glu  
35 40 45

Lys Leu Gly Trp Tyr His Ser Met Glu Ala Val Arg Arg Ala Phe Arg  
50 55 60

50 Val Trp Glu Gln Ala Thr Pro Leu Val Phe Gln Glu Val Pro Tyr Glu  
65 70 75 80

55 Asp Ile Arg Leu Arg Arg Gln Lys Glu Ala Asp Ile Met Val Leu Phe  
85 90 95

Ala Ser Gly Phe His Gly Asp Ser Ser Pro Phe Asp Gly Thr Gly Gly  
100 105 110

60 Phe Leu Ala His Ala Tyr Phe Pro Gly Pro Gly Leu Gly Gly Asp Thr  
115 120 125

His Phe Asp Ala Asp Glu Pro Trp Thr Phe Ser Ser Thr Asp Leu His  
130 135 140

	Gly Asn Asn Leu Phe Leu Val Ala Val His Glu Leu Gly His Ala Leu	145	150	155	160
5	Gly Leu Glu His Ser Ser Asn Pro Asn Ala Ile Met Ala Pro Phe Tyr	165	170	175	
	Gln Trp Lys Asp Val Asp Asn Phe Lys Leu Pro Glu Asp Asp Leu Arg	180	185	190	
10	Gly Ile Gln Gln Leu Tyr Gly Thr Pro Asp Gly Gln Pro Gln Pro Thr	195	200	205	
	Gln Pro Leu Pro Thr Val Thr Pro Arg Arg Pro Gly Arg Pro Asp His	210	215	220	
15	Arg Pro Pro Arg Pro Pro Gln Pro Pro Pro Pro Gly Gly Lys Pro Glu	225	230	235	240
	Arg Pro Pro Lys Pro Gly Pro Pro Val Gln Pro Arg Ala Thr Glu Arg	245	250	255	
	Pro Asp Gln Tyr Gly Pro Asn Ile Cys Asp Gly Asp Phe Asp Thr Val	260	265	270	
25	Ala Met Leu Arg Gly Glu Met Phe Val Phe Lys Gly Arg Trp Phe Trp	275	280	285	
	Arg Val Arg His Asn Arg Val Leu Asp Asn Tyr Pro Met Pro Ile Gly	290	295	300	
30	His Phe Trp Arg Gly Leu Pro Gly Asp Ile Ser Ala Ala Tyr Glu Arg	305	310	315	320
	Gln Asp Gly Arg Phe Val Phe Phe Lys Gly Asp Arg Tyr Trp Leu Phe	325	330	335	
	Arg Glu Ala Asn Leu Glu Pro Gly Tyr Pro Gln Pro Leu Thr Ser Tyr	340	345	350	
40	Gly Leu Gly Ile Pro Tyr Asp Arg Ile Asp Thr Ala Ile Trp Trp Glu	355	360	365	
	Pro Thr Gly His Thr Phe Phe Phe Gln Glu Asp Arg Tyr Trp Arg Phe	370	375	380	
45	Asn Glu Glu Thr Gln Arg Gly Asp Pro Gly Tyr Pro Lys Pro Ile Ser	385	390	395	400
	Val Trp Gln Gly Ile Pro Ala Ser Pro Lys Gly Ala Phe Leu Ser Asn	405	410	415	
	Asp Ala Ala Tyr Thr Tyr Phe Tyr Lys Gly Thr Lys Tyr Trp Lys Phe	420	425	430	
55	Asp Asn Glu Arg Leu Arg Met Glu Pro Gly Tyr Pro Lys Ser Ile Leu	435	440	445	
	Arg Asp Phe Met Gly Cys Gln Glu His Val Glu Pro Gly Pro Arg Trp	450	455	460	
60	Pro Asp Val Ala Arg Pro Pro Phe Asn Pro His Gly Gly Ala Glu Pro	465	470	475	480



Gly Ala Asp Ser Ala Glu Gly Asp Val Gly Asp Gly Asp Gly Asp Phe  
 485 490 495  
 5 Gly Ala Gly Val Asn Lys Asp Arg Gly Ser Arg Val Val Val Gln Met  
 500 505 510  
 Glu Glu Val Ala Arg Thr Val Asn Val Val Met Val Leu Val Pro Leu  
 515 520 525  
 10 Leu Leu Leu Leu Cys Val Leu Gly Leu Thr Tyr Ala Leu Val Gln Met  
 530 535 540  
 Gln Arg Lys Gly Ala Pro Arg Val Leu Leu Tyr Cys Lys Arg Ser Leu  
 545 550 555 560  
 15 Gln Glu Trp Val

## (2) INFORMATION FOR SEQ ID NO:9:

20

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 582 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: not relevant  
 25 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

30

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

35 Met Ser Pro Ala Pro Arg Pro Ser Arg Cys Leu Leu Leu Pro Leu Leu  
 1 5 10 15  
 Thr Leu Gly Thr Ala Leu Ala Ser Leu Gly Ser Ala Gln Ser Ser Ser  
 20 25 30  
 40 Phe Ser Pro Glu Ala Trp Leu Gln Gln Tyr Gly Tyr Leu Pro Pro Gly  
 35 40 45  
 Asp Leu Arg Thr His Thr Gln Arg Ser Pro Gln Ser Leu Ser Ala Ala  
 50 55 60  
 45 Ile Ala Ala Met Gln Lys Phe Tyr Gly Leu Gln Val Thr Gly Lys Ala  
 65 70 75 80  
 Asp Ala Asp Thr Met Lys Ala Met Arg Arg Pro Arg Cys Gly Val Pro  
 85 90 95  
 50 Asp Lys Phe Gly Ala Glu Ile Lys Ala Asn Val Arg Arg Lys Arg Tyr  
 100 105 110  
 55 Ala Ile Gln Gly Leu Lys Trp Gln His Asn Glu Ile Thr Phe Cys Ile  
 115 120 125  
 Gln Asn Tyr Thr Pro Lys Val Gly Glu Tyr Ala Thr Tyr Glu Ala Ile  
 130 135 140  
 60 Arg Lys Ala Phe Arg Val Trp Glu Ser Ala Thr Pro Leu Arg Phe Arg  
 145 150 155 160  
 Glu Val Pro Tyr Ala Tyr Ile Arg Glu Gly His Glu Lys Gln Ala Asp

	165	170	175
	Ile Met Ile Phe Phe Ala Glu Gly Phe His Gly Asp Ser Thr Pro Phe		
	180	185	190
5	Asp Gly Glu Gly Gly Phe Leu Ala His Ala Tyr Phe Pro Gly Pro Asn		
	195	200	205
10	Ile Gly Gly Asp Thr His Phe Asp Ser Ala Glu Pro Trp Thr Val Arg		
	210	215	220
	Asn Glu Asp Leu Asn Gly Asn Asp Ile Phe Leu Val Ala Val His Glu		
	225	230	235
15	Leu Gly His Ala Leu Gly Leu Glu His Ser Ser Asp Pro Ser Ala Ile		
	245	250	255
	Met Ala Pro Phe Tyr Gln Trp Met Asp Thr Glu Asn Phe Val Leu Pro		
	260	265	270
20	Asp Asp Asp Arg Arg Gly Ile Gln Gln Leu Tyr Gly Gly Glu Ser Gly		
	275	280	285
	Phe Pro Thr Lys Met Pro Pro Gln Pro Arg Thr Thr Ser Arg Pro Ser		
	290	295	300
25	Val Pro Asp Lys Pro Lys Asn Pro Thr Tyr Gly Pro Asn Ile Cys Asp		
	305	310	315
30	Gly Asn Phe Asp Thr Val Ala Met Leu Arg Gly Glu Met Phe Val Phe		
	325	330	335
	Lys Glu Arg Trp Phe Trp Arg Val Arg Asn Asn Gln Val Met Asp Gly		
	340	345	350
35	Tyr Pro Met Pro Ile Gly Gln Phe Trp Arg Gly Leu Pro Ala Ser Ile		
	355	360	365
	Asn Thr Ala Tyr Glu Arg Lys Asp Gly Lys Phe Val Phe Phe Lys Gly		
	370	375	380
40	Asp Lys His Trp Val Phe Asp Glu Ala Ser Leu Glu Pro Gly Tyr Pro		
	385	390	395
45	Lys His Ile Lys Glu Leu Gly Arg Gly Leu Pro Thr Asp Lys Ile Asp		
	405	410	415
	Ala Ala Leu Phe Trp Met Pro Asn Gly Lys Thr Tyr Phe Phe Arg Gly		
	420	425	430
50	Asn Lys Tyr Tyr Arg Phe Asn Glu Glu Leu Arg Ala Val Asp Ser Glu		
	435	440	445
	Tyr Pro Lys Asn Ile Lys Val Trp Glu Gly Ile Pro Glu Ser Pro Arg		
	450	455	460
55	Gly Ser Phe Met Gly Ser Asp Glu Val Phe Thr Tyr Phe Tyr Lys Gly		
	465	470	475
60	Asn Lys Tyr Trp Lys Phe Asn Asn Gln Lys Leu Lys Val Glu Pro Gly		
	485	490	495
	Tyr Pro Lys Ser Ala Leu Arg Asp Trp Met Gly Cys Pro Ser Gly Gly		
	500	505	510

Arg Pro Asp Glu Gly Thr Glu Glu Glu Thr Glu Val Ile Ile Ile Glu  
515 520 525

5 Val Asp Glu Glu Gly Gly Gly Ala Val Ser Ala Ala Ala Val Val Leu  
530 535 540

Pro Val Leu Leu Leu Leu Leu Val Leu Ala Val Gly Leu Ala Val Phe  
545 550 555 560

10 Phe Phe Arg Arg His Gly Thr Pro Arg Arg Leu Leu Tyr Cys Gln Arg  
565 570 575

Ser Leu Leu Asp Lys Val  
15 580

## (2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:  
20 (A) LENGTH: 607 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: not relevant  
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Ile Leu Leu Thr Phe Ser Thr Gly Arg Arg Leu Asp Phe Val His  
1 5 10 15

35 His Ser Gly Val Phe Phe Leu Gln Thr Leu Leu Trp Ile Leu Cys Ala  
20 25 30

Thr Val Cys Gly Thr Glu Gln Tyr Phe Asn Val Glu Val Trp Leu Gln  
35 40 45

40 Lys Tyr Gly Tyr Leu Pro Pro Thr Asp Pro Arg Met Ser Val Leu Arg  
50 55 60

45 Ser Ala Glu Thr Met Gln Ser Ala Leu Ala Ala Met Gln Gln Phe Tyr  
65 70 75 80

Gly Ile Asn Met Thr Gly Lys Val Asp Arg Asn Thr Ile Asp Trp Met  
85 90 95

50 Lys Lys Pro Arg Cys Gly Val Pro Asp Gln Thr Arg Gly Ser Ser Lys  
100 105 110

Phe His Ile Arg Arg Lys Arg Tyr Ala Leu Thr Gly Gln Lys Trp Gln  
115 120 125

55 His Lys His Ile Thr Tyr Ser Ile Lys Asn Val Thr Pro Lys Val Gly  
130 135 140

60 Asp Pro Glu Thr Arg Lys Ala Ile Arg Arg Ala Phe Asp Val Trp Gln  
145 150 155 160

Asn Val Thr Pro Leu Thr Phe Glu Glu Val Pro Tyr Ser Glu Leu Glu  
165 170 175

	Asn Gly Lys Arg Asp Val Asp Ile Thr Ile Ile Phe Ala Ser Gly Phe	180	185	190
5	His Gly Asp Ser Ser Pro Phe Asp Gly Glu Gly Gly Phe Leu Ala His	195	200	205
	Ala Tyr Phe Pro Gly Pro Gly Ile Gly Gly Asp Thr His Phe Asp Ser	210	215	220
10	Asp Glu Pro Trp Thr Leu Gly Asn Pro Asn His Asp Gly Asn Asp Leu	225	230	235
	Phe Leu Val Ala Val His Glu Leu Gly His Ala Leu Gly Leu Glu His	245	250	255
15	Ser Asn Asp Pro Thr Ala Ile Met Ala Pro Phe Tyr Gln Tyr Met Glu	260	265	270
20	Thr Asp Asn Phe Lys Leu Pro Asn Asp Asp Leu Gln Gly Ile Gln Lys	275	280	285
	Ile Tyr Gly Pro Pro Asp Lys Ile Pro Pro Pro Thr Arg Pro Leu Pro	290	295	300
25	Thr Val Pro Pro His Arg Ser Ile Pro Pro Ala Asp Pro Arg Lys Asn	305	310	315
	Asp Arg Pro Lys Pro Pro Arg Pro Pro Thr Gly Arg Pro Ser Tyr Pro	325	330	335
30	Gly Ala Lys Pro Asn Ile Cys Asp Gly Asn Phe Asn Thr Leu Ala Ile	340	345	350
	Leu Arg Arg Glu Met Phe Val Phe Lys Asp Gln Trp Phe Trp Arg Val	355	360	365
35	Arg Asn Asn Arg Val Met Asp Gly Tyr Pro Met Gln Ile Thr Tyr Phe	370	375	380
40	Trp Arg Gly Leu Pro Pro Ser Ile Asp Ala Val Tyr Glu Asn Ser Asp	385	390	395
	Gly Asn Phe Val Phe Phe Lys Gly Asn Lys Tyr Trp Val Phe Lys Asp	405	410	415
45	Thr Thr Leu Gln Pro Gly Tyr Pro His Asp Leu Ile Thr Leu Gly Ser	420	425	430
	Gly Ile Pro Pro His Gly Ile Asp Ser Ala Ile Trp Trp Glu Asp Val	435	440	445
50	Gly Lys Thr Tyr Phe Phe Lys Gly Asp Arg Tyr Trp Arg Tyr Ser Glu	450	455	460
55	Glu Met Lys Thr Met Asp Pro Gly Tyr Pro Lys Pro Ile Thr Val Trp	465	470	475
	Lys Gly Ile Pro Glu Ser Pro Gln Gly Ala Phe Val His Lys Glu Asn	485	490	495
60	Gly Phe Thr Tyr Phe Tyr Lys Gly Lys Glu Tyr Trp Lys Phe Asn Asn	500	505	510
	Gln Ile Leu Lys Val Glu Pro Gly Tyr Pro Arg Ser Ile Leu Lys Asp			

	515	520	525
5	Phe Met Gly Cys Asp Gly Pro Thr Asp Arg Val Lys Glu Gly His Ser 530 535 540		
	Pro Pro Asp Asp Val Asp Ile Val Ile Lys Leu Asp Asn Thr Ala Ser 545 550 555 560		
10	Thr Val Lys Ala Ile Ala Ile Val Ile Pro Cys Ile Leu Ala Leu Cys 565 570 575		
	Leu Leu Val Leu Val Tyr Thr Val Phe Gln Phe Lys Arg Lys Gly Thr 580 585 590		
15	Pro Arg His Ile Leu Tyr Cys Lys Arg Ser Met Gln Glu Trp Val 595 600 605		

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/26214

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/57 C12N9/64 C07K16/40 C07K19/00 C12N1/21  
C12N5/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

A	<p>PUENTE X S ET AL: "MOLECULAR CLONING OF A NOVEL MEMBRANE-TYPE MATRIX METALLPROTEINASE FROM A HUMAN BREAST CARCINOMA" CANCER RESEARCH, vol. 56, no. 5, 1 March 1996, pages 944-949, XP000644455 cited in the application</p>	
---	--	--

A	<p>MUELLER C G ET AL: "Polymerase chain reaction selects a novel disintegrin proteinase from CD40-activated germinal center dendritic cells." JOURNAL OF EXPERIMENTAL MEDICINE, (1997 AUG 29) 186 (5) 655-63. JOURNAL CODE: I2V. ISSN: 0022-1007., XP002101608 United States</p>	
---	--	--

-/--



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

29 April 1999

Date of mailing of the international search report

18/05/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Van der Schaal, C

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/26214

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>BERNOT A ET AL: "A transcriptional Map of the FMF region."            GENOMICS, (1998 JUN 1) 50 (2) 147-60.            JOURNAL CODE: GEN. ISSN: 0888-7543.,            XP002090815            United States            see table 2            -&amp; EMBL/GENBANK DATABASES Accession no            AJ003147, Sequences reference HSAJ03147            22 January 1998            XP002101614            see translation of nucleotides 6772-15161            -----</p>	1-8

